

RESPONSE OF SOME OLIVE CULTIVARS TO POTASSIUM LEVELS UNDER SALINITY CONDITIONS IN HYDROPONICS SYSTEM

Atta, Y. I. , A.F Hasan and A. M. Abdel-Fatah

Water Management Research Institute (NWRC), Egypt



ABSTRACT

A study was carried out for the effect of different levels of salinity i.e. (0,60 and 120mM (NaCl) and potassium levels i.e 0 and 30mM (K_2SO_4) on growth and chemical composition of seedlings of two olive cultivars karotina and picwal during 2009 and 2010 seasons at Inshas Research Station, El-Sharkia Governorate, Water Management Research Institute, Egypt. The seedlings were grown in hydroponics system (nutrient film technique NFT). Results revealed that, increasing salinity levels caused a gradually decrease in plant height, stem diameter, number of leaves/plant, number of branches/plant, N, P and K percentages, while Na, Ca and Mg percentages were increased gradually with increasing of salinity levels.

Potassium addition increased in plant height, stem diameter, number of leaves /plant, number of branches / plant, N, P and K percentages, while Na, Ca and Mg percentages were decreased. The results revealed the necessity of potassium fertilizer to reduce negative effects of salinity on plant growth.

Karotina cultivar gave the highest values of previous parameters than picwal cultivar and seems to be more than tolerate salinity compared with picwal cultivar.

Keywords: Salinity – Karotina – Sharkia – Hydroponice – Potasium

INTRODUCTION

Efforts have been done to develop hydroponics systems for using in propagation by cutting (Soffer and Burger (1988), Bertram (1991), and Tawfik (2001). These techniques have many advantages vis availability of high oxygen concentration which stimulate root initiation and development, facilitating of water influx to the cutting, simplicity and low cost.

In Egypt, wide areas of newly reclaimed lands are to be planted with different fruit trees using different water recourse such as from wells and drainage water. These resources may contain different levels of salinity. Fruit species, for example olive trees and even varieties exhibit different tolerance to level and kind of salinity (El-Saidi et al., 1988).

In most area where some fruit trees are irrigated with saline water enough for measurable effect on growth and yield, the main salts causing injury are chlorides (Abou - Rayya et al., 1988). Several researchers have investigated the effect of salinity on the growth of different species of mainly plants and they reported that, salinity conditions usually reduced plant growth according to type of salt and its concentration. In this respect. Downton (1977). explained this effect was a result of increasing level of chloride in the leaves which decreased the rates of CO_2 fixation. The reduction in photosynthesis could be attributed to increased residual (mesophyll) resistance to CO_2 fixation. El-Deen et al (1979). studied the effect increasing concentration of salinity levels from 0.0 to 4000 ppm on seedlings of olive (Shamlaliy cv) they found that plant height, number of leaves and leaf area index were reduced with sharply as a result to increase salinity levels. Patil and Waghmare (1983). who mentioned that all indices that were studied on pomegranate plants such as plant height, number of leaves, stem diameter, plant spread, leaf area decreased with increasing salinity. Hassan et al (1986). reported that olives appear to have medium salt tolerance. Hanafy (1989). on *Majarana hortensis* plants found that most salinity levels caused a clear reduction

in N,K and Ca percentages while Sodium percentage increased progressively as NaCl levels increased. Shehata and El-Tantawy (1994). on *Melia azederach* stated that the high levels of salinity reduced the fresh and dry weight of plants. El- Said et al (1995) studies on the susceptibility of six olive cultivars to salinity, they found that increasing of concentration of salts led to a decrease in plant height, stem diameter, leaves number/plant leaf area and total fresh and dry weight. They added that arrangement cultivars in tolerance to salinity were Meshan, kornaki, Tofahi, Manzanelo, Agizi, picwal, Ferdal, Hamedd, respectively. Azza and El-Mesiry (2000). on *Sesbania aegyptiaca* stated that increasing the salinity levels caused reduction in N and P contents while increased K content levels (250,500,1000 and 1500ppm). El-Ward et al (2004). Studied the effect of different salinity on growth of olive seedlings cv. "Khodiery". They found that the highest values of plant height, number of leaves/plant, number of branches/plant, stem diameter, length of root, fresh weight and dry weight of leaves and roots produced from control treatment in the two seasons while the lowest values of all previous characters produced from irrigation with saline water at the high level (1500ppm) and all the previous characters gradually decreased with increasing salinity concentration in the two seasons in most cases. Abd-Rabou, Farida et al (2008). Reported that increasing of concentration of salts led to a decrease in plant height, stem diameter, leaves number/plant, leaf area and total fresh and dry weight on Jackfruit and papaya seedlings.

Potassium (K^+) is considered to be one of the osmoregulating elements in plant cell especially in sugar storing plants. Frantzeskakis et al (1967) found that the potassium suppling power of highly calcareous soils was low when exchangeable K was less than 100 ppm under these conditions; it is highly probable that K deficiency symptoms will occur in olive trees. Rains and Epstein (1965) stated that potassium which is an essential cytoplasmic element, because of its involvement in osmotic regulation and it competitive effect against Na^+ . Yassoglou and Gavalas (1979) found a significant positive correlation between exchangeable

soil K and leaf K of Olive under nonsalin conditions. Awad and Rehm (1985). Found that application of high level of potassium significant increased the dry matter yield and some mineral contents of purple nutsedge plant. Kandil (1993). Suggested the increased K fertilization may be required under salinity conditions to contract the competitive effects of high concentrations of cations such as Ca^{2+} and Na^+ . Youssef and Saleh (1994). reported that K^+ is Known to counter act the harmful effects of Na. the increase in Na^+ and Cl^- and decrease in K^+ in different tissues of bean plants under salinity can be directly related to the reduction in the dry matter. Also salinity caused a reduction in total dry weight which was associated with increasing levels of sodium and chloride in plants and decreasing level of potassium uptake. And they added that the necessity of potassium fertilizer to reduce negative effects of salinity on plant growth. Many researchers reported that irrigation with saline sea water had a clear effect on decreasing plant growth. Hassan (1995). Reported that the under saline irrigation water use condition K deficiency systems can be disappeared through salinity hazard on leaves which led to dry out of the edges of leaves. Kandeil and El-Maddah (1997). reported that potassium addition generally increased the dry weight of leaves, stems and roots in bean plant at moderate rate (2000ppm) than the control one. And they found that potassium supply increased its content of been plant-Also, sodium content has increased up to application of 300ppm K-level and then decreased.

The objective of this work was to determine the effect of different levels of salinity as well as potassium levels on some growth and chemical content of two olive cultivars in hydroponics system.

MATERIALS AND METHODS

This investigation was conducted during the two successive seasons, 2009&2010 at Enshas Research Station, of Water Requirements Research El- Sharkia Governorate, Water Management and Irrigation

Systems Research Institute (National Water Research Center), which is located in East Nile Delta region in Egypt. The site is located at 30°-24' N. and 31 ° -35' E. longitude of with an elevation of about 25.5 meters above sea level.

The experiments were performed to study the effect of different salinity treatments as a form (NaCl) with or without addition of potassium element as a form Potassium Sulphate (K_2SO_4) under hydroponics system (nutrient film technique NFT). on the growth and chemical composition of some olive cultivars under lath-house conditions.

Hydroponics units were designed by using 6 plastic pipes. Each one pipe with 8 inches diameter and 6 m long with upper 18 holes every 30 cm with 10 cm diameter for putting olive seedlings were covered with a sheet of plastic for protective from direct of sunrays.

The pipes were putting up carriers with height 1 m made of galvanized iron with a gentle slope 1 cm/1m so that solution could be flowed under the influence of gravity. Each pipe was contact with tank of plastic contained 20 liters of solution. Each tank is equipped with small electrical pump for solution pumped in P.V.C hose (0.5 inch diameter) from the tank to higher end of pipe on shape film. Solution was flowed to lower end of pipe and accumulated in the tank with continuous recirculation.

The solutions were continuously aerated and were maintained at a pH of 6.5-7 by means of Hcl or NaoH. The solution was renewed twice-weekly, in order to avoid a significant reduction in concentration of nutrients.

Standard nutrient solution (S.N.S)

Standard nutrient solution (C.F. Douglas 1987) was used. It was component of commercial fertilizers (Macroelements) g/100 liter water and microelements mg/100 liter water as shown in Table (1) to given the following concentration of essential nutrient elements for seedlings of olive as a shown in Table (2).

Table (1). Amounts of Macro. and Microelements for Standard Nutrient Solution (S.N.S)

No	Macronutrient elements	g/100L water	No	Micronutrient elements	mg/100 L water
1	Urea $Co(NH_2)_2$	30	1	Iron Sulphate ($FeSO_4$).	6000
2	Calcium super phosphate $Ca(H_2PO_4)_2$.	40	2	Magnesium Sulphate ($MgSO_4$).	600
3	Potassium Sulphate (K_2SO_4).	60	3	Copper Sulphate ($CuSO_4$).	40
4	Calcium Nitrate ($CaNO_3$).	59	4	Zinc Sulphate ($ZnSO_4$).	40
5	Magnesium Sulphate ($MgSO_4$).	36	5	Boric acid (H_3BO_3).	180
6	Calcium Sulphate ($CaSO_4 \cdot 2H_2O$).	20	6	Ammonium Molubedate ($(NH_4)_6MO_7O_2$.	40
	Total weight	245		Total weight	6900

Table (2). Concentrations of Essential Nutrient Elements for Seedlings of Olive.

No	Element	mM/L	mg/L	No	Element	mM/L	mg/L
1	Nitrogen (N)	14.9	208	7	Iron (Fe)	0.218	12
2	Phosphorus (P)	2.4	75	8	Manganese (Mn)	0.036	2
3	Sulpher (S)	6.6	211	9	Zinc (Zn)	0.002	0.1
4	Calcium (Ca)	4.4	176	10	Copper (Cu)	0.002	0.1
5	Potassium (K)	7.5	294	11	Boron (B)	0.028	0.3
6	Magnesium (Mg)	2.0	50	12	Molybdenum (Mo)	0.002	0.2

Table (3): Effect of salinity and potassium levels on seedling height, seedling diameter No. of leaves /seedling and No. of branches/seedling of two olive cultivars in the two growing seasons and combined.

	seedling height (cm)			seedling diameter (mm)			No. of leaves /seedling			No. of branches/seedling		
	2009	2010	combined	2009	2010	combined	2009	2010	combined	2009	2010	combined
Salinity levels (S):												
S ₁ : Control	84.06	76.80	80.43	9.16	8.78	8.97	94.65	93.83	94.24	2.38	2.24	2.31
S ₂ : 60mM (NaCl).	80.55	73.63	77.09	7.04	6.82	6.93	75.73	73.75	74.74	1.89	1.72	1.81
S ₃ : 120mM (NaCl).	76.00	70.55	73.28	5.63	5.29	5.46	60.26	57.96	59.11	1.45	1.34	1.40
LSD 0.05	2.15	2.80	2.96	0.12	0.25	0.42	3.12	2.35	3.60	0.06	0.29	0.23
Potassium levels(K):												
K ₁ : 0 mM (K ₂ SO ₄).	79.36	72.88	76.12	7.13	6.85	6.99	75.23	73.93	74.58	1.82	1.64	1.73
K ₂ : 30mM (K ₂ SO ₄).	81.05	74.43	77.74	7.42	7.08	7.25	78.33	76.43	77.38	1.98	1.89	1.94
LSD 0.05	1.53	1.42	1.47	0.17	0.23	0.35	1.86	2.28	2.60	0.07	0.27	0.24
Cultivars (C):												
C ₁ : Picwal	77.05	70.98	74.02	6.95	6.61	6.78	73.32	71.88	72.60	1.82	1.67	1.75
C ₂ : Karotina	83.35	76.33	79.84	7.60	7.32	7.46	80.24	78.48	79.36	1.98	1.85	1.92
LSD 0.05	4.15	4.80	4.74	0.25	0.33	0.38	1.48	2.38	3.25	0.13	0.16	0.21
Interactions:												
SK	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S
SC	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S
KC	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S
SKC	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S	N.S

Also treating the seedlings with 30mM (K₂ SO₄) had thicker than the control in the two seasons and combined. These results agreed with those obtained by Thalooth et al (1990) who reported that potassium fertilization increased photosynthetic pigments content in sugar beet plant leaves. The positive effect of K⁺ for increasing chlorophyll content in plant leaves may be due to the indirect role for K⁺ element. In this respect pissark (1973) reported K-deficiency resulted in a collapse of chloroplasts in spring rape leaves.

Concerning the differences among cultivars regarding the stem diameter reached the significant level in two seasons and combined Table (3). The karotina cultivar had the thickest stem diameter and out yielded picwal by 9.11% these results, are in harmony with obtained by El-Said et al (1995).

No. of leaves/seedling

Data in Table (3) clearly show that the all salinity levels caused a reduction in No. of leaves/seedling compared with the control in the two seasons and combined. These results were emphasized by many investigators such as El-Deen et al (1979), Shehata (1992) on cupressus semperivens and Eucalyptus camaldulensis, Shehata and El-Tantawy (1994) on Melia azederach and El-Ward et al (2004) on olive as all of them stated that the high levels of salinity reduced the No. of leaves/seedlings.

For the effect of K levels on No. of leaves/seedling Data in Table (3) show clearly that it increased with increasing K concentration. These results are in agreement with those reported by kandil (1993), Hassan (1995).

There were significant differences in No. of leaves /seedling among the studied cultivars in the two seasons and combined. Karotina cultivar had the highest No. of leaves / seedling and out yielded picwal cultivar by 9.30% as average of the two seasons. Similar results were obtained by Hassan et al (1986), El-Said et al (1995).

No. of branches/seedling

The results listed in Table (3) showed that the values of No. of branches were least due to the increase in salinity levels and the decrease was significantly in the two seasons and combined.

This reduction in No. of branches/seedling were 21.65, and 39.40% at S₂ and S₃ respectively compared with the control S₁ treatment as average of the two seasons. Similar results reported by El-Ward et al (2004). These results are in agreement with those findings of Pearson and Hayward (1957) they found that, irrigation of grapefruit trees with a mixture of NaCl and CaCl₂ caused defoliation and die back. Also, the plants differed in degree of leaf burn and defoliation according to the water and soil salinity.

Results in Table (3) showed the applied of K₂ (30mM K₂SO₄) increased the No. of brunches/seedling by 9.70% compared with control. K₁ treatment (0.0 mM K₂SO₄). as average in the two seasons.

Effect of interactions:

The interactions between Salinity treatments, potassium treatments and cultivars were not significant in both seasons and combined for all traits (Table 3).

2. Chemical Composition

Data in Table (4) show the effect of k. levels and salinity levels on N, P, K, Na, Ca and Mg contents in the leaves of two olive cultivars. It is evident that salinity levels caused an increase in N content in the leaves. A Clear effect on reduction in N, P and K percentages as compared with the control Plants. In this respect, Hassan *et al* (1970) found a decrease of P uptake with increasing salinity, Also the decrease of phosphate translocation with high chloride supply was observed with sunflower. Ashour *et al* (1970), kandiell and El-Maddah (1997).

This result may indicate that the change in the balance between K and Na in the tissues may affect nucleic acid synthesis (Conway and lipman 1964). At the same time Na, Ca and Mg percentages increased as salinity level increased in the two seasons and their

averages. The control plants gave the highest percentages of (N, P, K), while the lowest percentages of (N, P, K), were obtained from the plants treated with the highest salinity levels (120mM Na Cl) in the two seasons and their averages. The highest Na, Ca and Mg percentages resulted from the seedlings treated with highest salinity level (120mM Na Cl), while the lowest percentages Na, Ca and Mg percentages were obtained from control treatment in the two seasons and their averages.

These results were in agreement with those obtained by Hanafy (1989). on *Majorona hortensis* who stated that most salinity levels caused a clear reduction in N, K and Ca percentages while Na percentage increased progressively as NaCl levels increased in both seasons. These results are in agreement with those of Hanafy (1994) on Fennel. The obtained results were similar to those obtained by Salem et al (1998). on

Dodonaea viscosa, Azza and El-Mesiry (1999) on *leuccana leucocephala*, Azza and El-Mesiry (2000) on *Sesbania egyptiaca* and Azza (2001) on *Casuarina glauca*

As for K levels data presented in Table (4) indicated that, there were a gradual increase in the percentages of N, P and K ions in the olive leaves with increased K level up to 30mM (K₂ SO₄), while decreased the percentages of Na, Ca and Mg ions. The obtained results are parallel with the finding of Rains and Epstein (1965), Frantzekakis et al (1967), Yassoglou and Gavalas (1979), Awad and Rehm (1985), Kandel (1993"), Youssef and Saleh (1994), Hassan (1995) and Kandilel and El-Maddah (1997). The contents of the elements N, P, K, Na, Ca and Mg in the cultivars of olive as affected by potassium and salinity treatments presented in Table (4).

Table (4): Effect of Salinity X K- Levels on Chemical Composition of Dry Leaves of Two Olive Cultivars (Average of Two Growing Seasons).

Minerals Content in leaves		N%			P%			K%			Na%			Ca%			Mg%		
Cultivars (C)	K- levels (mM/L) Salinity levels (mM/L)	K ₁	K ₂	Mean	K ₁	K ₂	Mean	K ₁	K ₂	Mean	K ₁	K ₂	Mean	K ₁	K ₂	Mean	K ₁	K ₂	Mean
		(0.0mM/L)	(30.0mM/L)		(0.0mM/L)	(30.0mM/L)		(0.0mM/L)	(30.0mM/L)		(0.0mM/L)	(30.0mM/L)		(0.0mM/L)	(30.0mM/L)		(0.0mM/L)	(30.0mM/L)	
Picwal C ₁	S1: control	3.65	3.80	3.73	0.16	0.21	0.19	1.35	1.52	1.44	0.53	0.48	0.51	0.42	0.38	0.40	0.24	0.21	0.23
	S2: 60mM (Na cl)	3.18	3.26	3.22	0.11	0.15	0.13	1.28	1.45	1.37	0.80	0.72	0.76	0.51	0.44	0.48	0.33	0.28	0.31
	S3: 120mM (Na cl)	2.71	2.96	2.84	0.09	0.12	0.11	1.15	1.37	1.26	0.98	0.86	0.92	0.67	0.52	0.60	0.36	0.33	0.35
	Mean	3.18	3.34	3.26	0.12	0.16	0.14	1.26	1.45	1.36	0.77	0.69	0.73	0.53	0.45	0.49	0.31	0.27	0.29
Karotina C ₂	S1: control	3.72	3.90	3.81	0.18	0.23	0.21	1.42	1.61	1.52	0.64	0.55	0.60	0.48	0.42	0.45	0.30	0.25	0.28
	S2: 60mM (Na cl)	3.21	3.33	3.27	0.15	0.19	0.17	1.37	1.53	1.45	0.88	0.80	0.84	0.55	0.51	0.53	0.41	0.33	0.37
	S3: 120mM (Nacl)	2.80	3.20	3.0	0.12	0.14	0.13	1.24	1.42	1.33	1.10	0.92	1.01	0.70	0.62	0.66	0.45	0.40	0.43
	Mean	3.24	3.48	3.360	0.15	0.19	0.17	1.34	1.52	1.43	0.87	0.76	0.82	0.58	0.52	0.55	0.39	0.33	0.36
Average	S1: control	3.69	3.85	3.77	0.17	0.22	0.20	1.39	1.57	1.48	0.59	0.52	0.56	0.45	0.40	0.43	0.27	0.23	0.25
	S2: 60mM (Na cl)	3.20	3.30	3.25	0.13	0.17	0.15	1.33	1.49	1.41	0.84	0.7m	0.80	0.53	0.48	0.51	0.37	0.31	0.34
	S3: 120mM (Na cl)	2.76	3.08	2.92	0.11	0.13	0.12	1.20	1.40	1.30	1.04	0.89	0.97	0.69	0.57	0.63	0.41	0.37	0.39
	Mean	3.22	3.41	-	0.14	0.17	-	1.31	1.49	-	0.82	0.mu	-	0.56	0.48	-	0.35	0.30	-

The data showed that the karotina cultivar had the highest percentages of N, P, K contents. Picwal cultivar had the highest Percentages of Na, Ca, and Mg content. In general there were different variations between potassium and salinity levels on N, P, K, Na, Ca and Mg contents of olive cultivars. These results are in agreement with those of El-Saidi *et al* (1988). And El-Ward et al (2004).

From the data illustrated in Tables (3 & 4), it can be concluded that Seedlings of two olive cultivars can tolerate chloride salt concentration up to 60mM, but increasing salt concentration to 120mM affected badly their growth and chemical properties, And consequently K application had reducing the injurious consequences of salinity on plant growth, whenever other measures are unavailable. The results revealed the necessity of potassium fertilizer to reduce negative effects of salinity on olive plant growth.

Karotina cultivar was more tolerance of salinity from picwal cultivar.

The work of the statistical analysis of the data using analysis of variance in three Ways and show off the results of statistical analysis proved a significant effect of salinity and potassium levels on the elements of dry matter in leaves of olive as proven moral statistical categories on the items in the dry article to olive leaf except for the component (N) did not demonstrate statistical moral his.

(LSD) to find differences between the salt and potassium and varieties levels have been using the separation test between the averages and the results were as described in Table (5) where it is clear that there are significant differences statistically between salinity and potassium levels on the elements of dry matter of the olive leaf as proven moral influence of varieties on the elements Article dry olive leaves except the element (N).

Table (5): The Results of Variance in Three Ways Analysis

		Sum of Square	Mean Square	DF	F	R ²
N	Cult	0.0300000	0.0300000	1	70.69**	0.97
	Salinity	1.4703500	0.73517500	2		
	Klevel	0.11603333	0.11603333	1		
B	Cult	0.00240833	0.00240833	1	59.34**	0.97
	Salinity	0.01211667	0.00605833	2		
	Klevel	0.00440833	0.00440833	1		
K	Cult	0.01840833	0.01840833	1	255.29**	0.99
	Salinity	0.06615000	0.03307500	2		
	Klevel	0.09900833	0.09900833	1		
NA	Cult	0.0225333	0.0225333	1	136.09**	0.98
	Salinity	0.3492666	1.1746333	2		
	Klevel	0.0300000	0.0300000	1		
Ca	Cult	0.00963333	0.00963333	1	44.65**	0.96
	Salinity	0.08351667	0.0417583	2		
	Klevel	0.01613333	0.1613333	1		
Mg	Cult	0.01267500	0.1267500	1	77.50**	0.97
	Salinity	0.03751667	0.01875833	2		
	Klevel	0.00700833	0.00700833	1		

CONCLUSIONS

In Egypt, wide areas of newly reclaimed lands are to be planted with different fruit trees using different water recourse such as from wells and drainage water. These resources may contain different levels of salinity. Fruit species, for example olive trees and even varieties exhibit different tolerance to level and kind of salinity.

Potassium (k⁺) is considered to be one of the osmoregulating elements in plant cell especially in sugar storing plants. The potassium suppling power of highly calcareous soils was low when exchangeable k was less than 100ppm under these conditions; it is highly probable that k deficiency symptoms will occur in olive trees.

This investigation was conducted during the two successive seasons, 2009&2010 at Enshas Research Station, of Water Requirements Research El- Sharkia Governorate, Water Management and Irrigation Systems Research Institute (National Water Research Center), which is located in East Nile Delta region in Egypt. The Seedlings were grown in hydroponics system (nutrient film technique NFT).

Results revealed that, increasing salinity levels caused a gradually decrease in plant height, stem diameter, number of leaves/plant, number of branches/plant, N, P and K percentages, while Na ,Ca and Mg percentages were increased gradually with increasing of salinity levels.

Potassium addition increased in plant height, stem diameter, number of leaves /plant, number of branches / plant, N, P and K percentages, while Na, Ca and Mg percentages were decreased. The results revealed the necessity of potassium fertilizer to reduce negative effects of salinity on plant growth.

Karotina cultivar gave the highest values of previous parameters than picwal cultivar and seems to be more than tolerate salinity compared with picwal cultivar.

REFERENCES

- Abd El-Aziz., Y.E.Y., M.M. Sharaf and El-Ashraf, M.A. (1985). Effect of Salinity, Sodium Adsorption Ratio and Chloride Levels on Vegetative Growth and Leaf Mineral Composition in Some Citrus Rootstock Seedling. 1-Effect on Vegetative Growth. *Annals of Agric. Sci. Moshtohor*, 23.
- Abd-Rabou, Farida A., Kassim F.S.E. and El-khaly, L.A. (2008). Effect of Water Salinity on Growth and Minerals Content of Jackfruit and Papaya Seedling. *Egypt. J. of Appl. Sci.*, Vol. 23 No. (6B) : 665 – 680.
- Abou – Rayya, M.S., Kasim N.E., and. El – Saidi M.T (1988). Effect of Irrigation with Saline Water on the Growth and Some Chemical Properties of Grape Transplants. *J. Agric. Sci., Mansouro Univ.* Vol. 13. No. (4) : 1634 – 1641.
- Allen, O.M. (1959). *Experiment in Soil Bacteriology Minneapolis, U.S.A Burgess Puble . Co.* p. 83-88.
- Ashour, N. I.; M. O. Kabesh and El-Oksh, I. (1970). Effect of chloride salinity on the absorption, exudation and distribution of phosphorus P³² by sunflower plants. *Agrochemical* 462.
- Awad, F. and. Rehm, S. R (1985). Effect of Replacing Potassium by Sodium on the Propagative Capacity and Mineral Content of Purple Nutsedge (Cyprus rotundas L.) in Full Light and Shade *Egypt. J. Soil Sci.*, 25: 49-61.
- Azza, A. Mazher and El-Mesiry, T. A. (1999). Effect of Salt on the Response of *Leucaena Leucocephala* to Biofertilization in Sandy Soils. *Recent Technologies in Agri Culture, Proc. of the 1st Congress. Cairo Univ. Fac. Of Agric.* 27-29 November Bull. Fac. Agric. Cairo Univ.

- Azza A.M. Mazher and El-Mesiry, T. A. (2000). Effect of Gamma Irridation on Growth and Some Chemical Constituents of Sesbania Aegyptiaca Seedlings Grown under Salt Stress Conditions. Egypt J. Appl. Sci. 15 (11): 110-116.
- Azza, A. Mazher (2001). Effect of Different Salinity Levels of diluted Sea Water and Soil Media on Growth and Chemical Composition of Casuarinas Glauca. Egypt Sci., Vol 16 No. (7): 26-37.
- Bertram, L. (1991). Vegetative propagation of Hibiscus rosasinensis L. in relation to nutrient concentration of the propagation medium, Science Horticulturae 48:131-139.
- Conway, T. W. and F. lipman (1964). Characterization of a Ribosome Linked Guanosine Triphosphatase in Escherichia coli Extracts. Proc. Natl. Acad. Sci, U.S.A. 52, 1469.
- Douglas, J.S. (1987). Advanced guide to hydroponics. Bralce, pub. Luc. New York. U.S.A.
- Downton, W.J.S. (1977). Photosynthesis in Salt Stressed Grapevines Division of Horticultural Research, CSIRO, G. P. O. Box 350 Adelaide, S.A. 5001. Australian Journal of Plant physiology 4(2):183-192.
- El-Deen S.A.; M.M. Wassif and Fadl, M.S. (1979). Effect of Salinity on Olive Seedling Growth. Agric. Res. Rev. 57:259-269.
- El – Said, M.E.; H.E. Emtithal, A. Hamoda and Sari. Eldin, S.A. (1995). Studies on the Susceptibility of Some Olive Cultivar to Salinity. Zagazig J. Agric Res., 22: 2314 – 2328.
- El-Saidi , M.T., N.E. kasim and M.S.M Abou –Rayya (1988). Effect of irrigation with salin water on the growth and some chemical proerties of picual olive seedlings . J. Agric. Sci. Mansoura Univ. Vol. 13 No. (4) : 1627 – 1633.
- El-Ward, H.D, M. S. Hanafy and A. M. El-Musawa (2004). Effect of Irrigation with Saline Water by Sodium Chloride and Calcium Chloride 1:1 on the Growth of Olive Seedlings CV Khodiery. Egypt. J. Appl. Sci. Vol 19 No. (11B): 623-634.
- Frantzeskakis, J., N. Yassoglou and J. Androulakis (1967). An Investigation of the Potassium Status in Some Olive. Gartenbauwiss Enschafft No. 42, pp. 113-117.
- Hanafy, M.S. (1989). Effect of Chemical Fertilization and Salinity on Production of Herb and Essential oil of Marjoram Plants (Majorana Hortensis Moench). ph. D. Thesis, Fac. Agric., Cairo Univ.
- Hanefy, M. S. (1994). Effect of Irrigation with Diluted Saline sea Water on Growth, yield, Essential Oil, productivity and Chemical Composition of Fennel (Foeniculum Vulgate L.) Plants. Egypt, J. Appl. Sci., Vol. 9 No. (8): 445-461.
- Hassan, M. A. M. (1995). Effect of Soil and Irrigation Water Properties on Potassium Content of Olive Leaves Grown in Delta of Wadi El-Arish . Zagazig J. Agric. Res. Vol. 22 No. (3): 943-953.
- Hassan, N. A. K., I. V. Drew, D. Knudsen and R. A. Olesen, (1970). Influence of Soil Salinity on production of dry matter and uptake and distribution of nutrients in barely and corn. Agrn. J. 62:43.
- Hassan M.M., A.A. Ibrahim, and M.A. Zahram (1986). Defferences in Salt Tolerance of Seedlings of Some Olive Cultivars. Egypt. J. Hort. 13:21-28.
- Jackson, M.L. (1962). Soil chemical Analysis Prentice – Hall, Inc. Englewood, Cliffs, N.7., U.S.A.
- Kandil A.A. (1993). Response of Some Sugar Beet Varieties to Potassic Fertilizers Under Salinity Conditions. Proc. of the ASWAS Conf., 8-15 Dec. 1990, Al Ain United Arab Emirates. Kluwer Academic Publishers, 199-207.
- Kandiel, SH. H. and El-Maddah, H. I. (1997). Effect of Salinity and Potassium Levels on Growth and Some Macronutrients Content of Snap bean Varieties Zagazig J. Agric-Res. Vol. 24 (1): 169-180.
- Patil, V.K. Waghmare, P.R. (1983). Tolerance Level of Exchangeable Sodium Percentage (ESP) for Pomegranate. Journal of Maharashtra Agricultural Univ. 8(3):257-259.
- Pearson, G. Z.A. and H. E. Hayward (1957). The Influence of Salinity and Water Table on Growth & Mineral Composition of Young Grapefruit Trees. Proc. Amer. Sco. Hort. Sci., 69: 197-203.
- Piper, C.S. (1950). Soil and Plant Analysis 158-750. The University of Adelaida, Adelaida.
- Pissark H. P. (1973). The Development of Potassium Deficiency Symptoms in Spring Rape. Z. Pflanzenernahr. Bodenk, 163: 1-96.
- Rains, D.W. and E. Epstein (1965). Transport of Sodium in Plant Tissues. Science (148): 1611.
- Salem M. A., A. A. Mazher and A. A. El-Malt (1998). Effect of Salinity Levels Soil Media and Gibberillic Acid Treatments on Hopseed Bush (Dodonaea Viscosa) Seedling Egypt J. Appl. Sc., Vol. 13 No. (9): 185-207.
- Saleh M. M., H. E. Wahba, and M. R. Farid (1993). Effect of Cycocel on the Resistance of Matricaria Chamomile L. to Salinity. Egypt. J. Appl. Sci, Vol 8 No. (11): 760-768.
- Sameni A. M. and Bassiri (1982). Interactive Effects of N, salinity Duration on Growth and Chemical Composition of broad bean. Iran. Agric. Res. 1(1): 49-60 (C. F. Faba bean Abst Vol. 3 No 4, 36 (1983).
- Shehata, M. S. (1992). Effect of Salinity and Soil Moisture Content on Seedling Growth of Suppresses Sempervirens L. and Eucalyptus Comaldulensis Dehn ph. D. Thesis, Fac. Agric Cairo Univ. Egypt.
- Shehata, M. S. and A. El-Tantawy (1994). Effect of GA₃ Spray on Growth and Some Chemical Constituents of Melia Azederach Seedlings Grown in Saline soil 1st Conference for Ornamental Horticulture, 22-24 October 1994, Fac. Agric., Cairo Univ.
- Snedecor, G.W. and W.G. Cochran (1974). Statistical Methods 6th ed. Iowa State Univ., Press, U.S.A.

- Soffer, H. and D. W. Burger (1988). Effects of Dissolved Oxygen Concentrations in aer-Hydroponics on the Formation and Growth of Adventitious Roots. J. Amer. Soc. Hort. Sci. 113 (2) 218-221.
- Tawfik, A. A. (2001) Fast and Efficient Mass-Propagation of Some Ornamental Plants Using Simple Water Air-Flow Medium. Acta Horticulture 548:611-618.
- Thalooth, A. T, H. A. El-Zeiny and A. O. M. Saad (1990). Application of Potassium Fertilizer for Increasing Salt Tolerance of Broad Bean. Bull of Egypt. Soc. Physiol. Sci, 10 (3): 182-198.
- Youssef, R.A. and Saleh, A.L. (1994). Effect of Salinity on Growth and Distribution of $Na^+ + K^+$ and Cl^- in bush bean Plants. Egypt.J. Appl. Sci., Vol., 9 No. (7): 691-697.
- Yassoglou N. J. and N. A. Gavalas, (1979). Soil and leaf Analysis of Olive Groves in Greece, Proc. 14th Coll. Int Potash. Inst. 279-288.

استجابة شتلات بعض أصناف الزيتون لمستويات البوتاسيوم تحت ظروف الملوحة في نظام المزارع المائية
يسرى إبراهيم عطا، على فرج حسن وأحمد محمد عبد الفتاح
معهد بحوث إدارة المياه – المركز القومي لبحوث المياه – مصر

أجرى هذا البحث في محطة بحوث المقننات المائية بإنشاص – محافظة الشرقية التابعة لمعهد بحوث إدارة المياه وطرق الري – المركز القومي لبحوث المياه خلال موسمى ٢٠٠٩ ، ٢٠١٠ لدراسة تأثير مستويات مختلفة من الملوحة (صفر ، ٦٠ ، ١٢٠ مللى مول من كلوريد الصوديوم (NaCl) ومستويات البوتاسيوم (صفر ، ٣٠ مللى مول من كبريتات البوتاسيوم (K_2SO_4) على بعض صفات النمو والمكونات الكيميائية على شتلات صنفين من الزيتون (بيكوال – كاروتينا).

وكانت أهم النتائج المتحصل عليها.

- زيادة مستويات الملوحة تسببت فى انخفاض تدرجى فى قيم كلاً من ارتفاع النبات ، قطر الساق ، عدد الأوراق /نبات ، عدد الأفرع/نبات وكذلك نسب K, P, N بينما زادت نسب كلاً من Mg , Ca , Na تدرجياً بزيادة مستوى الملوحة.
- زيادة مستويات البوتاسيوم أدت إلى زيادة قيم كلاً من ارتفاع النبات ، قطر الساق ، عدد الأوراق /نبات ، عدد الأفرع/نبات وكذلك نسب K, P, N بينما نقصت نسب كلاً من Mg , Ca , Na تدرجياً بزيادة مستوى الملوحة.
- ومن النتائج يمكن أن نستنتج أن إضافة البوتاسيوم ضرورى جداً لتقليل التأثير السلبى للملوحة على النمو والمكونات الكيميائية لصنفى الزيتون بيكوال – وكاروتينا محل الدراسة.
- أعطى الصنف كاروتينا أعلى القيم لصفات النمو السابقة وكذلك المكونات الكيميائية. هذا ويبدو أن الصنف كاروتينا أكثر تحمل لتأثيرات الملوحة عن الصنف بيكوال.