


Concentration of Zinc and Copper in Milk and Some Reproductive Aspects During Hot and Cold Seasons of Klebi Ewes Reared in South Egypt

Damarany, A. I.

Department of Animal and Poultry Production, Faculty of Agriculture and Natural Resources, Aswan University, Egypt.



ABSTRACT

The present study was carried out to evaluate concentrations of zinc and copper in milk, some reproductive aspects of Klebi ewes during the hot and cold seasons in south Egypt. According to season of lambing, 48 ewes lambed in the cold months, while 47 ewes lambed during hot months were used in this study. Results showed that concentrations of zinc and copper in Klebi ewe milk were higher (2.23 ± 0.3 and 0.48 ± 0.05 mg/l) in cold than in hot months (1.33 ± 0.2 and 0.2 ± 0.03 mg/l, $P < 0.05$), respectively. Lambing ($P \geq 0.05$) and twinning ($P < 0.05$) rates tended to be higher in cold than in hot months (122.9 and 106.4 and 37.3 and 12.0%, respectively). Stillbirth, abortion and retained placenta rates were more frequent in ewes lambed in hot than in cold months (8, 6, 4.3 vs. 5, 4, 2.1%, $P \geq 0.05$), respectively. Positive correlation coefficient ($P < 0.05$) between each of zinc or copper concentration in Klebi ewe milk and both lambing and twinning rates ($r = 0.615, 0.452, 0.631$ and 0.684 , respectively). Negative correlation coefficient ($P < 0.05$) was recorded between each of zinc and copper concentration in Klebi ewe milk and stillbirth, abortion and retained placenta rates. The current study demonstrated seasonal changes in concentration of zinc and copper in Klebi ewe milk, which were associated with pronounced alteration in reproductive process of Klebi ewes. Reproductive performance of Klebi ewes in cold was better than in hot season. Therefore the present study may conclude that dietary supplementation of zinc and copper, particularly during the hot season is recommended for improving reproductive aspects of Klebi ewes under environmental conditions of south Egypt.

Keywords: Klebi ewes, lambing season, milk, zinc, copper, reproduction.

INTRODUCTION

Sheep contribute to 1.3% from the total world milk production, according to FAO (2010). Majority of the sheep population in Egypt is in small holders (Ministry of agriculture, personal communication), most of the small holders depend on grazing for feeding their herds. Under extensive grazing system, the reproduction process of sheep was affected by nutrient availability, especially mineral contents of the forages. Machado *et al.* (2005) and Meeske *et al.* (2006) found fluctuation in total mineral content of ryegrass/clover pastures during different seasons. Michlova *et al.* (2016) reported that variability of the content of minerals in small ruminant milks related to the quality of feed and pasture, which is associated with fluctuations in vegetation and climate conditions.

The environmental conditions of south Egypt are characterized by widely seasonal variation, especially the ambient temperature (AT). The ambient temperature reached to 44.3 °C in hot and dropped sharply to 8.8 °C in cold months, particularly at night. This fluctuation in seasonal conditions reflected seriously on the quality and quantity of the pasture and the green fodders which consequently greatly reflected on the animal performance. The Klebi sheep is one of the local breeds that reared in south Egypt for meat production. Many investigations interested with the role of zinc and copper on the reproductive performance of sheep (Hemingway *et al.*, 2001; Vázquez-Armijo *et al.*, 2011; Abd-El-Monem *et al.*, 2015). The physiological functions of zinc is component of numerous metallo enzymes, influences transcription and cell replication and the deficiency lead to impaired spermatogenesis and development of secondary sex organs in males, reduced fertility in sheep and goats (Vázquez-Armijo *et al.*, 2011). While, copper is enzyme component and catalyst involved in steroidogenesis and prostaglandin

synthesis and its deficiency lead to delayed and depressed estrus, abortion, death fetuses, infertility, and congenital ataxia (Vázquez-Armijo *et al.*, 2011). Sales *et al.* (2011) reported that zinc and copper play main role in regulating progesterone production from corpus luteum via enzyme superoxide dismutase of crossbred heifers. Ceylan *et al.* (2008) reported that zinc effect on the secretion and function of testosterone by the enzymes that control the arachidonic acid. Zinc plays major role in the repair and maintenance of the uterine layer after parturition, which is necessary for embryo implantation (Hostetler *et al.*, 2003; Robinson *et al.*, 2006). Deficiency of copper in sheep diets lead to embryo loss, failures embryo implantation and fetal death (Hidiroglou, 1979).

Recently many investigation determined concentrations of minerals in milk of experimental animals as mirror of blood (Ranjith and Pandey, 2015; Farzad *et al.*, 2016; Michlova *et al.*, 2016). The present study aimed to spot light on concentration of zinc and copper in milk of Klebi sheep reared under traditional conditions (small holders and grazing system) and their relationships with some reproductive aspects under conditions of south Egypt.

MATERIALS AND METHODS

Location and climatic conditions:

This study was carried out in Aswan governorate; it is far from Cairo city by about 890 km. Sheep farm located in Kom Ombou city. The climatic conditions of Aswan governorate are famous for dry weather. The months of the year were divided into two seasons cold (November to April) and hot (May to October). Throughout the experimental period, ambient temperature, relative humidity and rainfall of cold and hot seasons were collected from Meteorological Authority and presented in Table (1).

Table 1. Ambient temperature ($^{\circ}$ C), relative humidity (%) and rainfall of cold and hot seasons throughout the experimental period.

Season	Ambient Temperature ($^{\circ}$ C)		Relative Humidity (%)	Rainfall mm./month
	Max.	Min.		
Hot	35.9 – 44.3	20.8 – 26.2	16 – 26	0.0 – 1.0
Cold	22.9 – 35.0	8.8 – 19.1	18 – 40	0.1 – 1

Animals and management:

A total number of 50 Klebi ewes were used in the present study. The age of animals ranged between 2 to 4 years and the parity ranged between 3rd and 5th. Live body weight of ewes ranged between 30 and 40 kg. Total of 48 ewes lambed in the cold months and 47 ewes lambed during the hot months. Animals were reared in traditional farm as semi-shaded yards. Ewes were fed on concentrate feed mixture (corn and wheat bran) at night and in the morning on residues of crops and plants for each season.

Reproductive aspects:

Reproductive parameters included: Rate of lambing, stillbirth, abortion, retained placenta and twinning) were recorded according to Tadege *et al.* (2015).

Lambing rate (%) = Number of lambs born/total number of lambed ewes. Stillbirth rate (%) = Number of lambs stillborn/total number of lambs born.

Abortion rate (%) = Number of ewes aborted/total number of pregnant ewes.

Retained placenta rate (%) = Number of ewes with retained placenta (≥ 12 hr)/number of lambed ewes.

Twinning rate (%) = Number of twins/total number of lambs born

Determination of zinc and copper:

Milk samples were collected from 48 ewes lambed in the cold months and 47 ewes lambed in hot months. Milk samples were taken at 6a.m. before drinking, after two weeks of post-partum period. The samples were analyzed quantitatively for zinc (Zn) and copper (Cu) based on the procedure outlined by the manufacture using atomic absorption (AA) flame (ICE 3000C113500040 v1.30, England) according to AOAC (2000).

Statistical analysis:

Statistical model included the effect of season of the year on different parameters by T. test using SAS (2002). Chi Square was used to compare between the percentage values, while person's correlation coefficients were calculated. The used model was:

$$Y_{ij} = \mu + B_i + e_{ij}$$

Where: Y_{ij} = the observations, μ = overall mean, B_i = effect of lambing season (cold =1, hot = 2) and e_{ij} = experimental error assumed to be randomly distributed (0, σ^2).

RESULTS AND DISCUSSION**Concentration of Zn and Cu in Klebi ewe milk:**

Overall mean of zinc and copper concentrations in Klebi sheep milk was 1.78 \pm 0.3 and

0.35 \pm 0.04 mg/l, respectively (Table, 2). The obtained value of zinc is within a range of 1.29 and 3.09 mg/l as reported by Khan *et al.* (2007) and Al-Wabel (2008), respectively. Meanwhile, the concentration of copper is within a range of 0.20 and 0.40 mg/l as reported by Barłowska *et al.* (2013) and El-Bagermi *et al.* (2014), respectively. The concentration of zinc and copper was lower than 24.14 and 0.63 mg/l (Michlova *et al.*, 2016), while was higher than 0.79 and 0.212 mg/l (Abed-Al-Helaly *et al.*, 2013), respectively.

Concentrations of zinc and copper in Klebi sheep milk are significantly ($P < 0.05$) higher in cold (2.23 \pm 0.3 and 0.48 \pm 0.05 mg/l) than in hot months (1.33 \pm 0.2 and 0.21 \pm 0.03 mg/l), respectively (Table 2). This result agreed with that reported by Khan *et al.* (2003), who found that concentration of zinc in sheep milk during the cold season was higher (1.29 mg/l) than in hot season (0.56 mg/l). In similar trend, Dar *et al.* (2014) found that concentrations of zinc and copper in ewe blood plasma in cold season was higher (1.17 and 1.38 mg/l) than in hot season (0.69 and 0.68 mg/l), respectively.

The present result is in agreement with that reported by Farzad *et al.* (2016), who found the grazing season was effective on variations of milk composition of mineral contents in Moghani sheep. They attributed the differences in milk mineral content to the soil and grazing plant. In addition, Haenlein (1996) reported that lactation, season, feeding and health of the animal cause difference in the composition of milk macro- or micro-elements. Suttle (2010) reported that some variables affect plant mineral composition such as plant genotype, soil environment and climate.

Table 2. Concentration (mg/l) of Zn and Cu in Klebi ewe milk during hot and cold months (mean \pm SE).

Element	Hot months	Cold months	Overall mean
Zn (mg/l)	1.33 \pm 0.212 ^a	2.23 \pm 0.343 ^b	1.78 \pm 0.323
Cu (mg/l)	0.21 \pm 0.032 ^a	0.48 \pm 0.052 ^b	0.35 \pm 0.041

^{a, b}: values within the same row having different superscripts are significantly different at ($P < 0.05$).

Effect of season of lambing on some reproductive aspects of Klebi ewes:

Lambing rate of Klebi ewes was 114.7% (Table 3). The present rate, was higher than that reported by Koyuncu and Yerlikaya (2007) and Aldomy *et al.* (2009), they found that the lambing rate ranged between 96.7 and 100%. Meanwhile the present values are lower than that reported by Teleb *et al.* (2009), being 150% in Saidi ewes. Lambing rate of Klebi ewes was higher in cold (122.9%) than in hot months (106.4%), but the differences were not significant (Table 3). The obtained result is in agreement with that reported by Teleb *et al.* (2009), who found that lambing rate was higher (160%) in winter than in summer season (140%) in Saidi ewes. Similar trend was reported by Koycegiz *et al.* (2009), who found that lambing rate during cold months was (127%) and in hot one (126%).

Percentage of stillbirth in Klebi ewes was 6.5% (Table, 3). These results are in agreement with the findings of Teleb *et al.* (2009), who reported that 6.1% of lambs were stillbirth in Saidi ewes. The obtained result is lower than that reported by Casellas *et al.* (2014), who found that stillbirth percentage was (9.6%) in lambs under semi-intensive system. Percentage of stillbirth of Klebi ewes was lower in cold (5%) than in hot months (8%), but the difference was not significant (Table 3). This result agreed with that reported by Teleb *et al.* (2009), who found that stillbirth was higher in summer (12.2) than in winter (1.5%) in Saidi ewes. Abortion rate of Klebi ewes was 5% (Table 3). The present result was lower than 7.5% as reported by Aldomy *et al.* (2009) and 7.6% (Zahraddeen *et al.*, 2010). Abortion rate was lower (4%) in cold than in hot months (6%), but the differences were not significant. In this respect, Miller *et al.* (1988) reported that deficiency of copper is one of the important factors responsible for early embryonic death. O'Donoghue and Boland (2002) found that lower zinc levels in the diet was associated with decreased in fertility and increase abortions in cattle. Percentage of retained placenta of Klebi ewes was 3.2% (Table 3). This result is lower than that reported by Hussain *et al.* (2013) and Alenyorege and Mensah (2015), who found the percentage of retained placenta in ewes ranged between 10.5 and 30.4%. Retained placenta percentage of Klebi ewes is higher (4.3%) in hot than in cold months (2.1%), while the differences was insignificant. The present result agreed with that reported by Alenyorege and Mensah (2015), who found that incidence of retained placenta was higher (36.3%) in hot than in cold months (34.5%). Campbell *et al.* (1999) reported that decrease in retained placentas in dairy cows fed diet supplemented with zinc and copper.

Twining rate of Klebi ewes was 25.7% (Table 3). The obtained result is higher than that reported by Aldomy *et al.* (2009) and Hussain *et al.* (2013), being 5 and 15%). Twining rate of Klebi ewes was significantly ($P < 0.05$) higher in cold (37.3%) than in hot months (12%). This result agreed with that reported by Teleb *et al.* (2009), who found twining rate in Saidi ewes was 63.6 and 40.8 in cold and hot season, respectively.

Table 3. Percentage of some reproductive aspects of Klebi ewes during hot and cold months.

Rate (%)	Hot months	Cold months	Overall mean
Lambing	106.4 (50/47)	122.9 (59/48)	114.7 (109/95)
Stillbirth	8 (4/50)	5 (3/59)	6.5 (7/109)
Abortion	6 (3/50)	4 (2/50)	5 (5/100)
Retained placenta	4.3 (2/47)	2.1 (1/48)	3.2 (3/95)
Twining	12 ^a (6/50)	37.3 ^b (22/59)	25.7 (28/109)

^{a,b}: values within the same row having different superscripts are significantly different at ($P < 0.05$).

The correlation between concentration of zinc in Klebi ewe milk and reproductive aspects:

The physiological functions of zinc is component of numerous metalloenzymes, influences transcription and cell replication and the deficiency lead to impaired spermatogenesis and development of secondary sex organs in males, reduced fertility in sheep and goats (Vázquez-Armijo *et al.*, 2011). Positive correlation

($P < 0.05$) was recorded between each of lambing and twinning rates and concentration of zinc in Klebi ewe milk. While negative correlation was observed between each of stillbirth, abortion and retained placenta rates and concentration of zinc in Klebi ewe milk (Table, 4).

The present results are in agreement with that reported by Vázquez-Armijo *et al.* (2011), who found that dietary Zn supplementation improved reproductive performance of ewes. Also, El-Nour *et al.* (2010) reported that conception rate of Baladi goats was increased from 60 to 80% after Zn treatment. In addition zinc plays major role in maintenance the epithelia layer of the reproductive organs, which is necessary for embryo implantation (Hostetler *et al.*, 2003; Robinson *et al.*, 2006). Zn deficiency affect the reproductive process in males and females like spermatogenesis, secondary sex organs development, estrus and gestation (Smith and Akinbamijo, 2000).

Table 4. Correlation coefficients between concentration of zinc in Klebi ewe milk and reproductive aspects.

Element	Reproductive aspect rates (%)				
	Lambing	stillbirth	Abortion	retained placenta	twining
Zn	0.615 [*]	-0.739 [*]	-0.327	-0.497	0.452

* values with having superscripts are significantly different at ($P < 0.05$).

The correlation between concentration of copper in Klebi ewe milk and reproductive aspects:

The physiological functions of copper is enzyme component and catalyst involved in steroidogenesis and prostaglandin synthesis and the deficiency lead to delayed and depressed estrus, abortion, death fetuses, infertility, congenital ataxia (Vázquez-Armijo *et al.*, 2011). Positive correlation coefficient ($P < 0.05$) was observed between each of lambing and twinning rates and concentration of copper in Klebi ewe milk. While negative correlation was observed between each of stillbirth, abortion and retained placenta and concentration of copper in Klebi ewe milk (Table, 5). The obtained results are in agreement with that reported by Abd El-Monem *et al.* (2015), who found that supplementation of copper to the diet of Baladi ewes improved estrus response, pregnancy, lambing and twinning rates. Vázquez-Armijo *et al.* (2011) found that supplementation of copper to diet of sheep affected reproductive process, like expression of estrus, embryo implantation and reduction in spermatogenesis. Ahmed *et al.* (2009) found that deficiency of copper has been associated with delayed estrus, low conception rates, infertility, and embryonic mortality in cattle and buffalo. Supplementation of copper and zinc to diets during spring and summer seasons improved the physiological status of sheep (Dar *et al.* 2014). Sales *et al.* (2011) reported that zinc and copper play main role in regulating progesterone production from corpus luteum via enzyme superoxide dismutase of crossbred heifers. Deficiency of copper in sheep diets lead to embryo loss, failures embryo implantation and fetal death (Hidiroglou, 1979).

Table 5. Correlation coefficient between concentration of copper in Klebi ewe milk and reproductive aspects.

Element	Reproductive aspects rates %				
	Lambing	Stillbirth	Abortion	Retained placenta	Twining
Cu	0.631*	-0.837*	-0.709*	-0.815*	0.684*

* values with having superscripts are significantly different at (P <0.05).

CONCLUSION

The current study demonstrated seasonal changes in concentration of zinc and copper in Klebi ewe milk, which were associated with pronounced alteration in reproductive process of Klebi ewes. Reproductive performance of Klebi ewes in cold was better than in hot season. Therefore the present study may conclude that dietary supplementation of zinc and copper, particularly during the hot season is recommended for improving reproductive aspects of Klebi ewes under environmental conditions of south Egypt.

REFERENCES

- Abd-El-Monem, U. M.; S.A. Peris and A. I.A. El-Shorbagy (2015). Effect of Copper sulphate supplementation on semen quality, ovarian activities and reproductive performance of Egyptian Baladi sheep. *J. Am. Sci.*,11(10):42- 50.
- Abed-Al-Helaly, L.; S. H. Rashed and L. F. Bdaiwi (2013). A Comparative Study of oxidant and antioxidant levels between human milk with other types of ruminant animals. *Iraqi Nat. J. Chem.*, 49: 86-99.
- Ahmed, W. M.; H. H. El Khadrawy;E. M. Hanafi; A. R. Abd El Hameed and H. A. Sabra (2009). Effect of copper deficiency on ovarian activity in Egyptian buffalo-cows. *World J. Zool.*, 4:1-8.
- Aldomy, F.; N. O. Hussein; L. Sawalha, K. Khatabeh and A. Aldomy (2009). A national survey of prenatal mortality in sheep and goats in Jordan. *Pakistan Vet. J.*, 29(3): 102-106.
- Alenyerege, B. and K. Mensah (2015). Incidence of retained placenta in ruminants and its treatment by rural farmers in northern Ghana. *Indian J. Appl. Res.*, 5 (7):434- 438.
- Al-Wabel, N. A. (2008). Mineral contents of milk of cattle, camels, goats and sheep in the central region of Saudi Arabia. *Asian J. Biochem.*, 3:373-375.
- AOAC (2000). Official Methods of Analysis. 17th Edition. Gaithersburg, Maryland, USA.
- Barłowska J.; A. Wolanciuk; M. Kędzierska-Matysek and Z. Litwińczuk (2013). Effect of production season on the basic chemical composition and content of macro- and microelements in cow and goat milk. *Żywn. Nauk. Technol. Jakość*, 6(91): 69-78.

- Campbell, M.H.; J.K. Miller and F.N. Schrick (1999). Effect of additional cobalt, copper, manganese, and zinc on reproduction and milk yield of lactating dairy cows receiving bovine somatotropin. *J. Dairy Sci.*, 82:1019-1025.
- Casellas, J.; G. Caja; X. Such and J. Piedrafita (2014). Survival analysis from birth to slaughter of Ripollesa lambs under semi-intensive management. *J. Anim. Sci.*, 85 (2): 512-517.
- Ceylan, A.; Y. Serin; H. Aksit and K. Seyrek (2008). Concentrations of some elements in dairy cows with reproductive disorders. *Bull. Vet. Inst. Pulawy.* 52: 109-112.
- El-Bagermi, M. A.; H. G. M. Edwards and A. I. Alajtal (2014). A comparative study on the physicochemical parameters and trace elements in raw milk samples collected from Misurata-Libya. *SOP Trans. Analy. Chem.*, 1 (2):15-23.
- Dar, A. A.; R. K. Jadhav; U. Dimri; A. A. Khan; H. M. Khan and M. C. Sharma (2014). Effects of physiological status and seasonal variation on plasma mineral profile of sheep in Kashmir valley. *Acad. J.*, 9 (4):69-76.
- El-Nour, H. M.M ; H. M. A. Abdel Rahman and S. A. El-Wakeel (2010). Effect of zinc-metthionine supplementation on reproductive performance , kids performance, minerals profile and milk quality in early lactating Baladi goats. *World Appl. Sci. J.*, 9 (3): 275- 282.
- FAO (2010). Statistics Division. Food and Agriculture Organization of the United Nations, Rome, Italy.
- Farzad M. A. Q.; R. V. Yonjalli; A. Ghorbani and B. F. Achachlouei (2016). Milk composition and mineral concentration affected by elevation and grazing season in the rangelands of north Sabalan mountain, Iran. *J. Adva. Agric. Tech.*, 3 (1):20-25.
- Haenlein, G.F.W. (1996). Nutritional value of dairy products of ewe and goat milk. *Int. Dairy Fed.*,19-21: 159-178.
- Hemingway, R.G; J. J. Parkins and N.S. Ritchie (2001). Enhanced reproductive performance of ewes given a sustained - release multi-trace element/vitamin ruminal bolus. *Small Rumin. Res.*, 39:25-30.
- Hidiroglou, M. (1979). Trace element deficiencies and fertility in ruminants: A review. *J. Dairy Sci.*, 62:1195-1206.
- Hostetler, C.E., R.L. Kincaid and M. A. Mirando (2003). The role of essential trace elements in embryonic and fetal development in livestock. *Vet. J.*,166:125-139.
- Hussain, S. O.; S. F. Al-Zubaidi and M. Asofi (2013). Different endometritis treatments in ewe: Comparative study. *J. Agric. and Vet. Sci.*, 3 (5): 91-94.
- Khan, Z. I.; M. Ashraf; M. Y. Ashraf; Z. Rahman and A. Hussain (2003). Mineral status of livestock (Goats and Sheep) based on soil , dietary components and animal tissue fluids in relation to seasonal changes and sampling period in specific

- region of Pakistan. J. Anim. Vet. Adva., 2: (8): 478- 495.
- Khan, Z. I.; M. Ashraf; I. Javed and S. E. Pollet (2007). Transfer of sodium from soil and forage to sheep and goats grazing in a semiarid region of Pakistan: Influence of the seasons. Trace Elem. Electrol., 24: 49-54.
- Koycegiz, F.; E. Emsen, C. A. G. Diaz and M. Kutluca (2009). Effects of lambing season, lamb breed and ewe parity on production traits of fat tailed sheep and their lambs. J. Anim. Vet. Adva., 8 (1):195-198.
- Koyuncu, M. and H. Yerlikaya (2007). Effect of selenium-vitamin E injections of ewes on reproduction and growth of their lambs. South Afric. J. Anim. Sci., 37 (3): 233-236.
- Machado, C.F.; T.S Morris ; J. Hodgson; and M. Fathalla (2005). Seasonal changes of herbage quality within a New Zealand beef cattle finishing pasture. N. Z. J. Agric. Res., 48 (2): 265-270.
- Meeske, R.; A. Rothauge; G. D. Van der Merwe and J. F. Greyling (2006). The effect of concentrate supplementation on the productivity of grazing Jersey cows on a pasture based system. South Afric. J. Anim. Sci., 36: 105-110.
- Michlova, T.; A. Hejtmanikova; H. Dragounova and S. Hornickova (2016). The content of minerals in milk of small ruminants. Agro. Res., 14: 1407-1418.
- Miller, J. K.; N. Ramsey and F. C. Madsen (1988). The Ruminant Animal. D.C. Church. Ed., Prentice Hall, Englewood Cliffs, N.J. USA, pp 342- 400.
- O'Donoghue, D.G. and M. Boland (2002). The effect of proteinated trace minerals on fertility and somatic cell counts of dairy cattle. J. Dairy Sci., 78: 248-255.
- Ranjith D. and J. K. Pandey (2015). Mineral Profiles in Blood and Milk of Sheep Inter. J. Sci. Res. 4 (10):821- 826.
- Robinson, J.J., C. J. Ashworth; J. A. Rooke; L. M. Mitchell ana T. G. McEvoy (2006). Nutrition and fertility in ruminant livestock. Anim. Feed. Sci. Tech., 126:259-276.
- Sales, J.N.S.; R.V.V. Pereira; R.C. Bicalho and P.S. Baruselli (2011) Effect of injectable copper, selenium, zinc and manganese on the pregnancy rate of crossbred heifers (Bos indicus × Bos taurus) synchronized for timed embryo transfer. Livestock Science. 142(1-3): 59-62.
- SAS (2002). User's Guide: Statistics, Version 9.0 Edition. SAS Institute Inc., Cary, NC, USA.
- Suttle, F. N. (2010) Mineral Nutrition of Livestock, CAB International, UK. 587 p.
- Tadeg, w. M.; F. R. Gudeta; T. y. Mekonen; Y. T. Asfaw; A. L. Birru and A. A. Reda (2015). Sero-Prevalence of small ruminant brucellosis and its effect on reproduction at Tellalak district of Afar region, Ethiopia. J Vet. Med. Anim. Health, 7(4):111-116.
- Teleb, D. F.; E. O. H. Saifelnasr and E. H. El-Sayed (2009). Factors affecting performance and survivability of Saidi lambs from lambing to weaning. Egyptian J. Sheep & Goat Sci., 4 (1): 55-74.
- Vázquez-Armijo, J. F.; R. Rojo; D. López; J. L. Tinoco; A. González; N. Pescador and I. A. Domínguez-Vara (2011). Trace Elements in Sheep and Goats Reproduction: A review. Trop. Subtrop. Agroecosy., 14: 1 – 13
- Zahraddeen, D.; I. S. R. Butswat and L.S. Taimako (2010). Assessment of reproductive problems in some ruminants under small holder husbandry system in Bauchi, Nigeria. J. Vet. Sci., 4: 1 – 8.

تركيزات الزنك والنحاس في اللبن وبعض المظاهر التناسلية خلال المواسم الحارة والباردة لنعاج الكلبى و المرباه في جنوب مصر

احمد إسماعيل ضمرائى

قسم الإنتاج الحيوانى، كلية الزراعة والموارد الطبيعية، جامعة أسوان، مصر

أجرى هذا البحث لدراسة تركيزات الزنك و النحاس فى لبن نعاج الكلبى و بعض المظاهر التناسلية للنعاج تحت الظروف البيئية فى جنوب مصر خلال مواسم السنة. تم تقسيم السنة إلى موسمين بارد و حار. بناء على موسم الولادة تم تقسيم النعاج الى مجموعتين الأولى هى مجموعة النعاج التى ولدت خلال الموسم البارد ، والثانية النعاج التى ولدت خلال الموسم الحار. أظهرت النتائج ان تركيزات الزنك والنحاس كان أعلى معنوياً فى عينات اللبن المأخوذة من نعاج الكلبى خلال الموسم البارد 2.23 ± 0.3 و 0.48 ± 0.05 مليجرام /لتر عن الموسم الحار 1.33 ± 0.2 و 0.03 ± 0.03 مليجرام /لتر على التوالي. كان معدل الولادات والتوائم أعلى فى النعاج التى ولدت خلال الموسم البارد عنه فى النعاج التى ولدت خلال الموسم الحار (122.9 ، 106.4 ، 18.6 و 6% على التوالي). سجل كلا من معدل نفوق المواليد بعد الولادة، والإجهاض و احتباس المشيمة اتجاهها معاكس حيث كانت هذه المقاييس أكثر تكراراً خلال الموسم الحار عنه فى الموسم البارد (8 ، 6 ، 4.3 و 4 ، 5 ، 4 ، 2.1% على التوالي). اتضح ان هناك معامل ارتباط معنوي موجب بين تركيز كل من الزنك والنحاس و معدل الولادات وتركيز النحاس و معدل التوائم (0.684 ، 0.631 ، 0.616 ، r). ايضاً لوحظ أن هناك معامل ارتباط معنوي سالب بين تركيزات النحاس فى عينات لبن النعاج وكل من معدل نفوق المواليد بعد الولادة، الإجهاض و احتباس المشيمة و كذلك بين تركيز الزنك و معدل نفوق المواليد. أظهرت الدراسة ان هناك تغيرات موسمية فى تركيزات الزنك والنحاس فى لبن نعاج الكلبى مرتبطة بتغيرات فى بعض المظاهر التناسلية. كان الأداء التناسلي لنعاج الكلبى التى ولدت خلال الموسم البارد أفضل من التى ولدت فى

Damarany, A.I.

الموسم الحار. وتوصى الدراسة بأهمية إضافة الزنك والنحاس إلى العلائق خلال الموسم الحار حيث انها قد تؤدي إلى تحسين بعض المظاهر التناسلية لنعاج الكلبى تحت الظروف البيئية لجنوب مصر .