

## **EFFECT OF CHAMOMILE FLOWERS MEAL AS FEED ADDITIVES IN LAYING JAPANESE QUAIL DIETS ON PRODUCTIVE AND REPRODUCTIVE PERFORMANCE**

**Abd El-Galil, K.\*; Henda A. Mahmoud\*; Amal M. Hassan\*\* and A.S. Morsy\*\***

**\* Animal and Poultry Nutrition Dept., Desert Research Center, El-Matariya, Cairo, Egypt.**

**\*\* Animal and Poultry Physiological Dept., Desert Research Center, El-Matariya, Cairo, Egypt.**

### **ABSTRACT**

A total number of 180 Japanese quail (120 females and 60 males) at 4 weeks of age were used in an experiment lasted 22 weeks. Experimental Japanese quail (*Coturnix coturnix japonica*) were divided randomly into four equal experimental groups (30 females in each group). The first group was fed a basal diet as control, while the other three groups were fed the a basal diet additives with the Chamomile flower meal (CFM), at levels of 0.25, 0.50 or 0.75 g/kg diet, respectively. The experimental diets were isocaloric (2900 kcal ME/kg), isonitrogenous (20% CP) and isofibrous.

The main objective of the present work was undertaken to establish the utilization of Chamomile flower (*Matricaria recutita L*) as medicinal plant feed additive in laying Japanese quail diets and its effect on productive and reproductive performance.

#### **Results obtained could be summarized as follows**

The final live body weight and body weight change during the whole experimental period increased significantly ( $P<0.05$ ) with increasing CFM levels. The highest final live body weight and body weight change were recorded by using 0.75 followed by 0.50 g/kg, while those fed control diet recorded the lowest ones.

Feed intake (g/day) increased significantly ( $P<0.05$ ) with increasing CFM levels, however 0.75 g/kg CFM recorded the highest one, while the control diet recorded the lowest ones.

The level of 0.50 g/kg CFM recorded the best ( $P<0.05$ ) feed conversion ratio (g feed /g egg mass), while the control diet recorded the worst ones

Age at sexual maturity and first egg weight recorded a non-significant difference among groups, while egg weight, egg number and egg mass during the whole experimental period recorded a significant difference ( $P<0.05$ ) among groups.

Hatchability and Fertility percentage recorded a significant difference among groups.

Egg yolk and shell thickness percentage showed a non-significant increase among the experimental groups, albumen and yolk index percentage were decreased ( $P<0.05$ ) by increasing CFM, while eggshell and egg shape were increased ( $P<0.05$ ) by increasing CFM as compared to the control group.

Digestibility coefficients of OM, CP, CF, EE, NFE and the nutritive values expressed as DCP, TDN % and ME (kcal/kg) were significantly varied ( $P<0.05$ ) among the different experimental groups. .

Level of 0.50 g/kg CFM group showed the best net revenue as well as the highest value of economic efficiency among experimental groups.

Level of 0.5 g/kg improved ( $P<0.05$ ) Sperm concentration, sperm motility, total motile sperm, live spermatozoa and semen quality factor.

Hydrogen ion concentration (pH) significantly decreased in level of 0.5 g/kg.

Cholesterol concentration decreased ( $P < 0.05$ ) in the Japanese quail fed with level of 0.5 and 0.75 g/kg.

Glucose, albumin, creatinine, ALT and AST concentration did not significantly as affected by using different levels of CFM.

**In conclusions**, It could be concluded that using dietary medicinal plant such as Chamomile flower (*Matricaria recutita L*) meal at 0.50 g/kg of the diet could improve productive, reproductive performance and economical efficiency of laying Japanese quail.

**Keywords:** *Quail, Chamomile meal (CFM), productive performance, egg quality, digestibility, economical efficiency and some physiological function.*

## INTRODUCTION

Many attempts have been made by poultry nutritionists to improve their productive performance and feed utilization in order to reduce the cost of feeding by using dietary additives such as antibiotics, probiotics, enzymes and herbal medicinal plants.

Recently, many countries tended to prohibit the using of antibiotics as growth promoters because of their side effect on both birds and human health.

Abd El-Galil (2007), Abaza, (2001), Gill (1999) Khodary *et al.* (1996) and Abdel-Aal and Attia (1993) indicated that addition of herbal parts and plant seeds as natural feed additives improved performance of poultry.

The recent studies have showed that medicinal plants can be used instead of chemical compounds in poultry diets as natural tonic, restoratives (Boulos, 1983), antibacterial and antiparasitic drugs (khodary *et al.*, 1996) to obtain the best performance parameters, immunity and the viability of birds (El- Hindawy *et al.*, 1996 and Osman, 1996).

The natural feed additives as the medicinal plants such as Chamomile flower which are used in the treatment of various diseases in human. This plant, not only serves as a medicinal purpose, but also contains aromatic substances and essential oils that used in food industries for human (Evans and pharm1975).

The genus, Chamomile (*Matricaria recutita L*) is an annual herb from the asteraceal compositae and native to the Mediterranean, Euro-Siberian, Irano-Siberian regions and cultivated in Germany, West Asia and Egypt

Egyptian chamomile famous by its high quality and therefore large quantities of this plant exported to west Europe quantities. The used parts are the dried flower heads. Cultivated area in Egypt is 5502 feddans /year of which 3927 feddans/year are found in Fayoum. The production quantity in Egypt is 4268 ton/year and in fayoum 2846 ton/year (A.E.P. 1994)

German chamomile (*Matricaria recutita L.*), one of the popular ingredients in herbal teas, has been traditionally used for medicinal purposes of human, Chamomile has moderate antioxidant and antimicrobial activities, and significant antiplatelet activity in vitro. Animal model studies indicate that potent anti-inflammatory action, some ant mutagenic and cholesterol-lowering activities, as well as antispasmodic and anxiolytic effects (McKay and Blumberg, 2009)

Chamomile contains coumarins, flavonoids and volatile oils (0.24-1.40%). Main component of volatile oils are (-) alpha-bisabolol (up to 50%) (Isaac 1997) and chamazulin (1-15%) (Mann and Staba,1986). The effective material of chamomile is Azulene, cynaroside, luteolin, umbelliferone and herniarin (Zeid, 1998).

So, the main objective of the present work was to establish the utilization of Chamomile flower as herbal feed additives in laying Japanese quail diets and its effect on reproductive performance, nutrient digestibility, economic efficiency and some physiological functions.

## **MATERIALS AND METHODS**

The present experiment was carried out at Maryiout Experimental Research Station (South West of Alexandria), which belongs to the Desert Research Center. A total number of 180 Japanese quail (120 females and 60 males) at 4 weeks of age were used in an experiment lasted 22 weeks. Experimental Japanese quail (*Coturnix coturnix japonica*) were kept under similar managerial, hygienic and environmental conditions and were divided randomly into four equal experimental groups of 30 females in each group.

Quail were kept in batteries, which were divided into separate cages, where two females were housed in each cage. The first group was fed the basal diet as control (0% CFM), while the other three groups were feed additive the Chamomile flower meal. Chamomile flower meal (CFM) was added to the control diet at level of 0.25, 0.50 or 0.75 g/kg, respectively.

The experimental diets (Table1) were formulated according to N.R.C (1994) and were isonitrogenous (20% crude protein) and isocaloric (2900 kcal ME/kg). Feed and water were available *ad libitum*. Chemical analysis of the experimental diet and dried excreta were assayed using methods of A.O.A.C (1990).

During the experimental period, individual live body weight and feed intake were determined biweekly. Feed conversion ratio (g feed intake / g egg mass) was calculated and the mortality was recorded every day.

Age at sexual maturity was determined by an average of the first ten eggs for each group.

Eggs were collected daily and weighed for each group, so egg number, egg mass were calculated during the experimental period. At 15 weeks of age, 20 eggs were randomly taken from each group and were used to evaluate egg quality; yolk. Weight and shell weight were recorded. Shell thickness (without membrane) was measured by micrometer, while albumen weight was calculated by subtracting yolk and shell weight from egg weight, Yolk, shell and albumen percentage were calculated as a percentage of egg weight.

Males were housed individually in cages (one quail per cage) and fed the same diets for females. At 15 weeks of age, males (40 males) were transferred to female cages (two females and one male) for twenty minutes

for five days, the eggs were then collected and incubated. Hatchability percentage was calculated for each group.

**Table (1): Composition and proximate chemical of basal diet.**

Ingredients	%
Yellow corn	61.23
Soybean meal (44% CP)	10.70
Concentrate (52% CP)*	10.00
Corn gluten meal (60% CP)	6.97
Wheat bran	5.47
Dicalcium phosphate	0.50
Limestone ground	4.45
Vit. and min. premix**	0.25
L-lysine	0.16
DL- methionine	0.27
<b>Total</b>	<b>100</b>
<b>Proximate chemical analysis %</b>	
Crude protein	20.29
Crude fiber	3.24
Ether extract	3.35
<b>Calculated values</b>	
Metabolizable energy (kcal/kg)***	2900
Calcium %	2.50
Available phosphorus %	0.30
Methionine %	0.45
Lysine %	1.00
Methionine + Cystin %	0.70
Price /kg diet (L.E)	1.593

\* Protein concentrate contained, 52 %Crude protein, 2.03% Crude fiber, 6.17% Ether extract, ME 2800 (kcal/kg), 1.50 % Methionine, 2.0% Methionine & Cystin, 3.0%Lysine 7.00% Calcium, 2.93 % Available Phosphorus 2.20 % Nacl.

\*\* Each 1 kg Vitamins and minerals premix contains, Vit. A 120000 IU, Vit. D3 20000 IU, Vit.E, 100 mg, Vit.K, 20mg, Vit. B1, 10 mg, Vit. B2, 50mg, Vit. B6,15 mg, Vit. B12, 100 µg, Pantothenic acid 100 mg, Niacin 300 mg, Folic acid 10mg, Biotin 500 IJg, Iron. 300mg, Manganese 600 mg, Choline chloride 500 mg., Iodine 10 mg, Copper 100 mg, Selenium 1 mg, Zinc 500 mg and 1200 mg Antioxidant.

\*\*\*\*Calculated according to price of feed ingredients at the same time (2008) of the experiment. Price of one kg chamomile flower meal =10.00 L.E.

At the end of the experimental feeding period, digestion trials were conducted using 20 adult quail males (five quail from each treatment) to determine the digestibility coefficients and the nutritive values of the experimental diets as affected by CFM levels. Males were housed individually in metabolic cages.

The digestibility trials extended for 9 days of them 5 days as a preliminary period followed by 4 days as collection period. The individual live body weights were recorded during the main collection period to determine any loss or gain in the live body weights. During the main period, excreta were collected daily and weighed dried at 60°C bulked finally ground and stored for chemical analysis. The faecal nitrogen was determined according to Jakobsen *et al.* (1960). Urinary organic matter was calculated according to

Abou-Raya and Galal (1971). Metabolizable energy was calculated according to Titus and Fritz (1971).

The digestion coefficients % of organic matter (OM), crude protein (CP), crude fiber (CF), ether extract (EE) and nitrogen free extract (NFE) of the experimental diets were estimated.

The nutritive values expressed as digestible crude protein (DCP), total digestible nutrients (TDN) and metabolizable energy (ME) were calculated.

Economical efficiency for egg production was calculated from the input / output analysis according to the costs of the experimental diets and selling price of one kg egg. The values of economical efficiency were calculated as the net revenue per unit of total costs.

Five quail from each treatment were chosen randomly for blood samples, Hemoglobin concentration was determined immediately colorimetric ally in fresh blood samples using readymade kits and the rest of the blood was centrifuged for 15 minutes at 3000 rpm to collect plasma before being stored at -20 °C until blood analysis.

The assays of plasma total protein, albumin, globulin (Value of globulin was calculated by subtracting the value of albumin from the value of total protein), glucose, cholesterol, creatinine, liver enzymes alanine transaminase (ALT) and aspartate transaminase (AST) were determined by the colorimetric methods using readymade kits.

Semen was collected from 5 sexually mature males from each treatment chosen randomly (at 15 weeks of age) by subjected the males to massages. The ejaculate volume was recorded by using a calibrated micropipette.

A dilute eosin solution (5%) was used for evaluating sperm concentration by the Neubauer haemocytometer slide according to Smith and Mayer (1955). Total sperm output was calculated by multiplying ejaculate volume and spermatozoa concentration.

Assessment of dead and abnormal spermatozoa was performed using an eosin–nigrosin blue staining mixture (Blom, 1950), then calculated as a percentage out of randomly chosen 100 sperm counted.

A drop of freshly collected semen was placed on a slide and the percentage of motile sperm was estimated by visual examination under low-power magnification (10 x) using a phase-contrast microscope according to Melrose and Laing (1970). Motility estimated according to Sedki (1999).

Semen quality factor (SQF) = (sperm concentration × ejaculate volume × live spermatozoa)/100.

Hydrogen ion concentration (pH) of semen was determined immediately after collection using pH paper.

Statistical analysis was carried out using General Linear Model (GLM) procedures by SAS program (1996) using simple one way analysis of variance according to this model:

$$Y_{ij} = \mu + T_i + e_{ij}$$

Where:

$Y_{ij}$  = Represented observation in  $j^{\text{th}}$  Chamomile flower meal

$\mu$ =Overall mean.

$T_j$  = Effect of  $j^{\text{th}}$  Chamomile flower meal ( $j = 0, 0.25, 0.5$  or  $0.75$  g).

$e_{ij}$  = Random error.

Duncan's New Multiple Range Test (Duncan, 1955) separated differences among treatment means.

## RESULTS AND DISCUSSION

### Live body weight and body weight change

Data for live body weight and body weight change during the whole experimental period of quail females is summarized in table (2). The final live body weight and body weight change during the whole experimental period varied significantly ( $P < 0.05$ ) among the experimental groups.

**Table (2): The productive performance ( $\bar{X} \pm \text{SE}$ ) of laying quail as affected by Chamomile flower meal addition.**

Items	Levels of Chamomile flower meal (g/kg)				Sig
	Control (0)	0.25	0.50	0.75	
Initial live body weight(g)	117.53+3.32	114.99+3.20	116.91+3.05	116.58+4.01	ns
Final live body weight (g)	247.78 <sup>b</sup> +5.91	250.01 <sup>ab</sup> +5.71	255.63 <sup>ab</sup> +4.51	257.36 <sup>a</sup> +6.08	*
change of live body weight (g)	130.25 <sup>b</sup> +2.26	135.02 <sup>ab</sup> +2.71	138.72 <sup>a</sup> +2.52	140.78 <sup>a</sup> +3.32	*
Age at sexual maturity/bird/day	49.22 +0.30	49.01+0.27	48.79+0.29	48.51+0.38	ns
First egg weight (g)	11.17 +0.20	11.14 +0.19	11.10+0.22	10.59+0.25	ns
Egg weight (g)	11.65 <sup>b</sup> +0.09	12.33 <sup>a</sup> +0.07	12.29 <sup>ab</sup> +0.05	12.00 <sup>ab</sup> +0.11	*
Egg number/ bird/ day	0.67 <sup>b</sup> +0.06	0.67 <sup>b</sup> +0.05	0.68 <sup>ab</sup> +0.03	0.69 <sup>a</sup> +0.05	*
Egg mass (g)/ bird/day	7.81 <sup>b</sup> +0.08	8.26 <sup>ab</sup> +0.02	8.36 <sup>a</sup> +0.03	8.28 <sup>ab</sup> +0.05	*
Feed intake (g)/ bird/day	29.21 <sup>b</sup> +0.11	29.64 <sup>ab</sup> +0.08	29.88 <sup>ab</sup> +0.07	30.10 <sup>a</sup> +0.10	*
Feed conversion ratio	3.70 <sup>a</sup> +0.06	3.59 <sup>ab</sup> +0.05	3.57 <sup>b</sup> +0.02	3.64 <sup>ab</sup> +0.03	*
Mortality rate %.	0.00	0.00	0.00	0.00	
Hatchability %	77.09 <sup>b</sup> +3.82	78.68 <sup>ab</sup> +2.16	79.89 <sup>a</sup> +2.01	79.10 <sup>a</sup> +3.11	*
Fertility %	89.43 <sup>b</sup> +4.36	91.39 <sup>ab</sup> +3.05	93.76 <sup>a</sup> +3.15	92.04 <sup>a</sup> + 4.07	*

<sup>a,b</sup>: Means within a row with different superscripts are significantly different.

Sig.= Significance, \* = ( $P < 0.05$ ), n.s = not significant.

It is worthy noting that final live body weight was improved with increasing the CFM level in the diet. Birds, gradually fed diets additive with 0.75 g/kg CM recorded 3.87 % higher than that of the control group, while 0.25 or 0.50 g/kg CM resulted in 0.90% and 3.17% higher than that of the control group, respectively. It is clear that live body weight change during the whole experimental period was increased by increasing CFM levels in the experimental diets. The addition with 0.25, 0.5 or 0.75 g/kg CFM level increased live body weight change by 3.66, 6.50 and 8.08% more than that of the control group, respectively.

The increase in body weight change may be due to the increase in feed intake and the improvement in nutrients digestibility of diets. Reflection on the positive effect may be attributed to the biological function of medicinal plant components that have been essential for growth (Boulos, 1983; Bradley, 1992 and Leung and Foster, 1996). Ibrahim *et al.* (1998) reported that adding natural additives to broiler diets caused a significant increase in broiler performance.

#### **Age at sexual maturity**

Sexual maturity age ranged from 49.22 to 48.51day, showing that CFM in laying quail diets did not affect this trait as shown in table(2).

#### **Weight of first egg**

Results of first egg weight recorded a non significant decrease among groups. It was noticed that there was a decrease in weight of first egg with the increase of CFM in the diets. It is worthy noting that feeding quail on 0.75 g/kg.CFM recorded the lowest values that may be attributed to early sexual maturity compared to other experimental groups (table 2).

#### **Egg weight**

Egg weight during the whole experimental period significantly differ ( $P<0.05$ ) among the experimental groups. It is worth noting that addition of diet by 0.25, 0.50 or 0.75 g/kg of CFM recorded an increase in egg weight amounted to 5.84, 5.49or3.00% compared to the control group, respectively.

Level of 0.25 g/kg recorded the highest egg weight (12.33g) while Level of 0.75 g/kg CFM recorded the lowest egg weight (12.00g), compared to other treated groups of CFM level as shown in table (2).

#### **Egg number and egg mass**

Results in (table 2) indicate that egg number and egg mass (Em) during the whole experimental period significantly varied ( $P<0.05$ ) among the experimental groups. It is worthy noting that egg number was higher in birds receiving 0.50 and 0.75 % CFM compared to other experimental groups, It is clear that addition of diet by 0.50 or 0.75 g/kg CFM increased egg number by1.49 or2.99 % than that of the control group, respectively. This may be due to the differences in earlier sexual maturity and attributed to the biological function of medicinal plant components. Khodary *et al.*(1996) stated that the efficiency of medicinal plants and some plant seeds as natural tonic, restoratives, antibacterial and anti-parasitic drugs in improving the productive performance in poultry.

Egg mass recorded maximum values for 0.50g/kg of CFM diet, while minimum values recoded for 0.25 g/kg CFM level.

It is worthy noting that feeding quail on 0.25, 0.50 or 0.75 g/kg of CFM resulted in 5.80, 7.00 and 6.00 % higher in egg mass than that of the control group, respectively.

The increase in egg mass with the 0.50 or 0.75 g/kg of CFM level was expected in view of the increase in egg number and vice versa with the 0.25 g/kg of CFM level.

#### **Feed intake and feed conversion ratio**

On the basis of the whole experimental period there was a significant differences ( $P<0.05$ ) among the experimental groups for daily feed intake trait. The experimental groups exhibited more feed intake compared to the control group. It is clear that increasing CFM levels in the experimental diets increased that feed intake. The addition with 0.25, 0.50 or 0.75 g/kg of CFM increased feed intake by1.13, 2.29 and 3.18 % more than that of the control group, respectively.

Natural feed additives had beneficial effect for stimulation and activity of digestive system by improving the diet palatability and enhancing appetite of poultry, thus increasing the amount of feed consumed (Namur et al., 1988).

The results of feed conversion ration showed significant improvement in feed efficiency utilization by the CFM level in diet as compared to the control group during the experimental period. Quail fed 0.50 g/kg CFM during the experimental period recorded the best feed conversion. Fritz et al., (1992) suggested that Chamomile improve feed conversion of poultry.

The data indicated that the feed conversion ratio of the group fed diets added with CFM was improved compared to those of the control group.

The improvement in feed conversion ratio of 0.50 g/kg CFM may be due to its highest egg mass as compared to that of CFM levels. Such improvement may be attributed to the properties of these materials that could act not only as antibacterial, anti-protozoa and antifungal but also as antioxidants (Bradley, 1992 and Leung and Foster, 1996).

#### **Mortality**

No incidence of mortality occurred during the experimental period as well as no effects of CFM levels supplemented on the experimental diets (Table 2). Eisenberg *et al.* (1993) indicated that medicinal plant contains natural substances that can promote health and alleviate illness.

#### **Hatchability**

Results of hatchability showed significant ( $P < 0.05$ ) improvement by the CFM levels in diet as compared to the control group (Table 2).


However, feeding quail on 0.25, 0.50 or 0.75% CFM resulted in 1.54, 3.10 and 2.08 % higher in hatchability than that of the control group, respectively.

Level of 0.50 g/kg recorded the highest hatchability compared to other groups of CFM level which may be due to the increase in fertility and improvement of ( $P < 0.05$ ) Sperm concentration compared to other experimental groups.

#### **Fertility %**

Results on fertility percentage in the present study recorded a significant difference among groups (Table 2).

It is worthy noting that feeding quail on 0.50 g/kg CFM level recorded higher values. This may be attributed to an improvement of sperm concentration, sperm motility, total motile sperm, live spermatozoa and semen quality with level of 0.5 g/kg, (Fig.1).



### Egg quality traits

The comparison between diets containing different levels of CFM and egg quality in comparison with the control diet (Table 3).

Data on yolk and shell thickness percentage showed a non-significant increase among the experimental groups, while albumen and yolk index percentage were decreased ( $P < 0.05$ ) by increasing CFM as compared to the control group. Eggshell and egg shape were significantly ( $P < 0.05$ ) increased by increasing CFM as compared to the control group.

**Table (3): Egg quality ( $\bar{X} \pm SE$ ) as affected by Chamomile flower meal in laying quail diets.**

Items	Levels of Chamomile flower meal (g/kg)				Sig.
	Control (0)	0.25	0.50	0.75	
Egg weight (g)	11.50+0.11	11.57+0.09	11.63+0.08	11.61+0.10	n.s
Yolk %	31.11 <sup>a</sup> +0.08	31.35 <sup>ab</sup> +0.09	31.50 <sup>a</sup> +0.06	31.66 <sup>a</sup> +0.07	n.s
Albumen %	55.12 <sup>a</sup> +0.09	55.10 <sup>ab</sup> +0.07	54.42 <sup>ab</sup> +0.04	54.11 <sup>b</sup> +0.07	*
Egg shell %	13.20 <sup>b</sup> +0.03	13.04 <sup>ab</sup> +0.04	13.71 <sup>a</sup> +0.04	14.01 <sup>a</sup> +0.05	*
yolk index %	48.70 <sup>a</sup> +0.10	48.39 <sup>a</sup> +0.09	47.57 <sup>ab</sup> +0.05	46.96 <sup>b</sup> +0.09	*
Egg shape %	78.14 <sup>b</sup> +0.05	78.50 <sup>ab</sup> +0.02	79.02 <sup>ab</sup> +0.04	81.22 <sup>a</sup> +0.03	*
Shell thickness (mm)	0.240+0.03	0.240+0.04	0.241+0.02	0.241+0.05	n.s

<sup>a,b</sup>: Means within a row with different superscripts are significantly different.

Sig.=Significance, \*= ( $P < 0.05$ ), n.s = not significant.

### Digestibility and nutritive values

Apparent digestion coefficients values of dietary treatments are illustrated in table(4), regarding those of organic matter (OM), crude protein (CP), crude fiber (CF), ether extract (EE) and nitrogen free extract (NFE).

Such values were significantly ( $P < 0.05$ ) differed among the experimental groups and the data indicated that, all nutrients digestibility values increased for quail fed CFM diet compared to the control diets.

**Table (4): Digestibility coefficients and nutritive values ( $\bar{X} \pm SE$ ) of the experimental diets as affected by Chamomile flower meal.**

Items	Levels of Chamomile flower meal (g/kg)				Sig
	Control (0)	0.25	0.50	0.75	
<b>Digestion coefficients %</b>					
OM	79.05 <sup>b</sup> +1.62	79.84 <sup>ab</sup> +1.32	80.02 <sup>ab</sup> +1.40	81.75 <sup>a</sup> +1.39	*
CP	80.51 <sup>b</sup> +1.25	81.62 <sup>ab</sup> +1.18	82.19 <sup>a</sup> +1.20	82.99 <sup>a</sup> +1.24	*
CF	24.08 <sup>b</sup> +1.51	24.78 <sup>b</sup> +1.20	26.29 <sup>ab</sup> +1.24	27.49 <sup>a</sup> +1.11	*
EE	86.98 <sup>b</sup> +1.22	87.95 <sup>ab</sup> +0.90	88.29 <sup>ab</sup> +0.82	89.05 <sup>a</sup> +0.71	*
NFE	86.95 <sup>b</sup> +1.25	88.95 <sup>a</sup> +1.13	88.97 <sup>a</sup> +1.20	89.57 <sup>a</sup> +1.26	*
<b>Nutritive values</b>					
DCP%	16.33 <sup>b</sup> +0.30	16.56 <sup>ab</sup> +0.21	16.68 <sup>ab</sup> +0.28	16.83 <sup>a</sup> +0.33	*
TDN%	64.16 <sup>b</sup> +1.31	65.42 <sup>ab</sup> +1.30	65.62 <sup>ab</sup> +1.35	66.15 <sup>a</sup> +1.41	*
ME(kcal/kg)	2701 <sup>b</sup> +26.16	2754 <sup>b</sup> +22.03	2761 <sup>ab</sup> +25.08	2785 <sup>a</sup> +37.70	*

<sup>a,b</sup>: Means within the same row showing different letters are significantly different. Sig.=Significance, \*( $P < 0.05$ ).

Generally, there was a significant increase in the digestion coefficients of OM, CF, EE and NFE of the experimental diets when compared with the control diet.

Similar trend was observed for the nutritive values of the tested diets in terms of digestible crude protein (DCP %), total digestible nutrients (TDN %) and ME (kcal/kg), which significantly ( $P < 0.05$ ) increased with the increasing of CFM up to 0.75 g/kg level.

It is important to note that the results of the digestion trials were coincided generally with the differences in production, productive performance and feed conversion ratio in quail diets in comparison with the control diet.

These results are in agreement with Abd El-Galil (2007) and Abd EL-latif *et al*, (2003) who indicated that, addition of medicinal herbal plants had a significant effect on improving digestibility coefficient and nutritive values.

Adding herbal medicinal plants to the diet asserted the biological role for herbal medicinal plants in activities of metabolic functions and biosynthesis of hormones (Abd El-Latif *et al.*, 2002).

#### **Economical efficiency**

The present results indicated that the diet containing 0.50 g/kg CFM as a feed additive result in the highest net revenue and economic efficiency compared to the other experimental groups as shown in (table 5). This may be related to the improvement of feed conversion ratio.

It was noticed that the control diet gave the lowest net revenue and economic efficiency. The best relative economic efficiency was detected with 0.50 g/kg CFM being 106.53% followed by those fed 0.25 g/kg CFM, respectively when compared with the control group.

**Table (5): Economical evaluation of quail as affected by Chamomile flower meal.**

Items	Control	Levels of Chamomile flower meal (g/kg)		
		0.25	0.50	0.75
Feed conversion ratio	3.70	3.59	3.57	3.64
Cost of kg feed (L.E)	1.593	1.593	1.594	1.594
Feed cost of kg egg (L.E)	5.894	5.719	5.691	5.802
Selling price of one kg egg (L.E)	13.00	13.00	13.00	13.00
Net revenue (L.E)	7.106	7.281	7.309	7.198
Economic efficiency	120.56	127.31	128.43	124.06
Relative economic efficiency%	100	105.60	106.53	102.90

**Semen characteristics**

Mean values of the Japanese quail semen characteristics as affected by different levels of Chamomile flower meal (CFM) are presented in table (6).

**Table (6): Effect of Chamomile flower meal as feed additives on the semen characteristics of Japanese quail.**

Items	Control (0)	Levels of Chamomile flower meal (g/kg)			Sig.
		0.25	0.50	0.75	
Ejaculate volume(ml)	0.018±0.001	0.018±0.001	0.020±0.002	0.023±0.003	n.s
Sperm conc. (×10 <sup>6</sup> ml)	663.33 <sup>b</sup> ±31.7	816.66 <sup>a</sup> ±52.3	836.66 <sup>a</sup> ±46.6	703.33 <sup>ab</sup> ±8.8	*
Total sperm output (×10 <sup>6</sup> )	11.93±1.29	14.69±2.20	16.73±2.34	16.17±2.30	n.s
Sperm motility %	80.00 <sup>b</sup> ±5.77	86.66 <sup>ab</sup> ±3.33	96.66 <sup>a</sup> ±3.33	76.66 <sup>b</sup> ±3.33	*
Total motile sperm (×10 <sup>6</sup> )	9.54 <sup>b</sup> ±1.76	12.73 <sup>ab</sup> ±2.34	16.17 <sup>a</sup> ±2.56	12.39 <sup>ab</sup> ±1.15	*
Live spermatozoa %	81.00 <sup>b</sup> ±3.78	85.33 <sup>ab</sup> ±2.33	91.33 <sup>a</sup> ±1.76	81.66 <sup>a</sup> ±5.84	*
Dead spermatozoa %	19.00 <sup>b</sup> ±3.78	14.67 <sup>ab</sup> ±2.33	8.67 <sup>a</sup> ±1.76	18.34 <sup>a</sup> ±5.84	*
Sperm abnormalities %	10.33±1.85	8.00±3.51	7.66±0.88	9.33±2.84	n.s
Semen quality factor	9.66 <sup>b</sup> ±1.53	12.53 <sup>ab</sup> ±2.11	15.27 <sup>a</sup> ±1.83	13.20 <sup>ab</sup> ±0.80	*
Hydrogen ion (pH)	7.83 <sup>a</sup> ±0.16	7.33 <sup>ab</sup> ±0.16	7.16 <sup>b</sup> ±0.16	7.66 <sup>ab</sup> ±0.16	*

a, b Means bearing different superscripts within the same row are significantly different (P<0.05).

Data revealed no significant differences in the ejaculate volume, total sperm output and sperm abnormalities between the treatment groups.

Sperm concentration was significantly increased (P<0.05) in the quail fed diet containing 0.50 g/kg CFM by 23.1% followed by 0.75 g/kg as compared to the control group.

Additionally, feeding Japanese quail with level of 0.5 g/kg improved (P<0.05) sperm motility, total motile sperm, live spermatozoa and semen quality factor by 20.8, 69.4, 12.7 and 58.0 %, respectively when compared to the control group. On the other hand, hydrogen ion concentration (pH) significantly decreased with level of 0.5 g/kg by 8.5 % as compared to the control group. White (1976) found that seminal plasma is usually an isotonic neutral medium and is detrimental to sperm cell survival. So, pH is a good indicator for semen quality (Morsy, 2007).

These improves in semen characteristics may attribute to the protective effect of CFM on the testis and epididymis and/or to the biological function of Chamomile flower meal as antioxidant and a scavenger of oxygen free radicals in seminal plasma which are toxic by-products of many metabolic processes (McKay and Blumberg, 2009). The improvement may be due acceleration of the mechanism of the reproductive hormones in the body. This effect is also apparent ( $P<0.05$ ) as improved fertility level by CFM additive in this experiment (Table 2). Khaki *et al.* (2009), they reported that the natural antioxidants can protect DNA and other molecules from cell damage induced by oxidation and can improve sperm quality and increase reproductive efficiency in rats.

**Hematological parameters**

Hematological parameters are presented in Table(7). Treating Japanese quail with CFM at a level 0.5 g/kg increased ( $P<0.05$ ) hemoglobin, total protein and globulin concentrations by 68.0, 45.7 and 58.7%, respectively, as compared to the control group. These findings are inagreement with those reported by El-Elamei (2001) in rabbits, Abaza (2001) in broiler, Nadia (2003) in broiler and Abu-Taleb *et al.* (2008) in Japanese quail. They all reported that using medicinal herbs such as chamomile improved digestion, limiting the development of undesirable microorganisms and promotion of bile flow facilitating digestion and absorption of nutrients. Hence, the improvement in hemoglobin may be due to increased iron absorption caused by CFM.

**Table (7): Effect of Chamomile flower meal as feed additives on the hematological parameters of Japanese quail.**

Items	Control (0)	Levels of Chamomile flower meal (g/kg)			Sig
		0.25	0.50	0.75	
Hemoglobin (g/dl)	10.75 <sup>b</sup> ±0.79	15.57 <sup>ab</sup> ±2.81	18.06 <sup>a</sup> ±0.28	11.33 <sup>b</sup> ±1.81	*
Glucose (mg/dl)	372.71±1.42	362.63±41.05	343.40±16.30	347.25±3.96	n.s
Total protein (g/dl)	6.45 <sup>b</sup> ±0.49	8.15 <sup>ab</sup> ±0.71	9.40 <sup>a</sup> ±0.48	8.49 <sup>a</sup> ±0.34	*
Albumin (g/dl)	3.35±0.71	3.50±0.61	4.48±0.49	4.40±0.09	n.s
Globulin (g/dl)	3.10 <sup>b</sup> ±0.54	4.65 <sup>a</sup> ±0.29	4.92 <sup>a</sup> ±0.28	4.09 <sup>ab</sup> ±0.40	*
Cholesterol (mg/dl)	135.40 <sup>a</sup> ±5.4	121.66 <sup>ab</sup> ±2.0	112.06 <sup>b</sup> ±10.33	107.39 ±5.8	*
Creatinine (mg/dl)	0.41±0.02	0.25±0.06	0.39±0.02	0.37±0.14	n.s
Alanine transaminase (ALT) (I.U./L)	30.66±8.25	30.00±5.79	21.00±5.50	20.66±5.23	n.s
Aspartate transaminase (AST) (I.U./L)	44.92±11.69	46.85±11.07	46.40±7.03	46.05±2.95	n.s

a, b Means bearing different superscripts within the same row are significantly different ( $P<0.05$ ).

The increase in total protein with CFM level of 0.5 g/kg by 45.7 % as compared to the control may be attributed to the improvement digestibility coefficient and nutritive values (Table 4). These findings are inagreement with those reported by Abd El-Galil (2007).

Cholesterol concentration decreased ( $P<0.05$ ) by 17.2 and 20.6 % with level of 0.5 and 0.75 g/kg, respectively, as compared to the control group. This result is in agreement with those obtained by Nadia (2003), Riyadh

(1987) and Sharma (1986). Abaza (2001) whom reported that feeding broiler chickens on diets containing 0.25 % chamomile flower significantly decreased total cholesterol by about 30.5% than the control group.

Glucose, albumin, creatinine, ALT and AST Concentrations did not significantly affected by using different levels of CFM.

The results showed that the addition of CFM up to 0.5 g/kg to Japanese quail diet had insignificant decrease of glucose level by -7.8 % compared to the control group. Such effect of CFM in lowering plasma glucose could be explained on the basis of increasing insulin binding sites of erthrocytes (Nadia, 2003).

The insignificant decrease in creatinine, ALT and AST concentrations may be due to the medicinal effect of CFM plant on the reductions in the severity of histopathological degeneration in liver. Also, this may indicates no harmful effect of Chamomile flower on liver and kidney tissues (Nadia,2003 and Schelicher *et al.*,1998).

In this respect, Adding herbal medicinal plants to the diet asserted the biological role for herbal medicinal plants in activities metabolic functions and biosynthesis of hormones (Abd El-Latif *et al.*, 2002). These effects may be due to the reflection of the antimicrobial, ant-fungal, antiseptic activities and anti-inflammatory (Abaza, 2001).

**In conclusions**, It could be concluded that using dietary medicinal plant such as Chamomile flower (*Matricaria recutita L*) meal at 0.50 g/kg of the diet could improve productive, reproductive performance and economical efficiency of laying Japanese quail.

## REFERENCES

- Abaza, I. M. (2001). The use of some medical plants as feed additives in broiler diets Ph.D. thesis, Fac. Agric. Alex. Univ. Egypt.
- Abd El-Galil, K., (2007). Effect of some feed additives on performance of laying Japanese quail. 1-Effect of Margoram leaves meal as a feed additive in laying Japanese quail diets. J. Agric. Sci. Mansoura Univ., 32(7):5167-5179.
- Abd El-Latif, S .A.; A.T.El-Yamany and Eman A. F. Edaly (2003). Evaluation of using different levels and sources of medical herbs in growing Japanese quail diets. Egypt.J.Nutr.and Feeds 6 (Specical Issue):219-220.
- Abd El-Latif, S.A.; Faten, A. A. Ibrahim and A. M. El-Kaiaty (2002).Effect of feeding dietary thyme black cumin,dianthus and fennel on peroductive and some metabolic responses of growing japanese quail. Egypt. Poult. Sci., 22(1): 109-125.
- Abd El-Latif, S .A.; A.T.El-Yamany and Eman A. F. El-Daly (2004). Evaluation of using different levels and sources of medical herbs in growing Japanese quail diets. Egypt.J.Nutr.and Feeds 7 (1):69-81.

- Abdel-Aal, E.S.M. and R.S. Attia (1993). Characterization of black cumin (*Nigella sativa*) seed. 1-Chemical composition and lipids. Alexandria of Science Exchange, 14:467-482.
- Abou-Raya, A.K. and A.G.H. Galal (1971). Evaluation of poultry feeds in digestion trials with reference to some factors involved. Egypt. J. of Anim., 11(1): 207-221.
- Abu Taleb, A.M., Hamodi, S.J. and El Afifi, Sh.F. (2008). Effect of using some medicinal plants (anise, chamomile and ginger) on productive and physiological performance of Japanese quail. Isotope and Radiation Research.
- A.E.P. (1994) Agricultural Economic Publication. Ministry of Agricultural and land reclamation. A.R.E.
- A.O.A.C. (1990). In "Official Methods of Analysis", Association of Official Agricultural Chemists, 15<sup>th</sup>. ed., Washington, D.C. USA.
- Blom, E. (1950). A 1-min live-dead sperm stain by means of eosin-nigrosin. *J. Fertil. Steril.*, 1: 176-177.
- Boulos, I. (1983). Medicinal plants of north Africa reference publication. Inc. Proc., 2nd Con. Fac. Vet. Med. Cairo, Univ., Cairo, Egypt, (1992).
- Bradley, P.R. (1992). British Herbal Compendium, Vol.1, Bournemouth: British Herbal Medicine Association.
- Duncan, D. B. (1955). The multiple range and F- tests. Biometrics, 11: 1-42.
- El-Elamei, Randa, R. E. (2001). Evaluation of some medicinal plants in rabbit nutrition. M.Sc. Thesis. Faculty of Agriculture, Cairo University.
- El-Hindawy, M.M; M.L. Tawfeek and S.H. Osman (1996). Effect of some biological feed additives for different periods on performance of broiler chick. Zagazig J. Agri. Res, Vol. 23 (5): 773-783.
- Eisenberg, D. M.; R.C. Kessler; C. Foster ; F.E. Norlock.; D. R., Calkins and T. L. Delbanco (1993). Unconventional medicine in the United States. Preference, costs and patterns of use. N. Engl J. Med, 328:246-252.
- Evans, F.J. and B. pharm (1975). Herbs, A concise guide in colour herbs by FERTILIZER. Frantisek severa, English consultant F.J. Evans, B. Pharm. Phosphorus. D. Hamlyn. London, New York. SYndeny, Toronto.
- Fritz, Z; A. Schleicher; S. Kinal; L. Tarosz and F. Majdanski (1992) Substitution of antibiotics by herbs in feed mixtures for broiler chicks Rozavway, Naukowe, Zootechniki, Moogrofie, Rozarway Poland, NO.31, 315-325.
- Gill, C. (1999). More Science behind 'Botanical: herbs and plant extracts as growth enhancers. Feed international (4): 20-23.
- Ibrahim, M. R.; M. S. Abd El-latif and A. T. El-Yamany, (1998). Effect of adding some natural growth promoters to broiler chicks diets on growth performance, digestibility and some metabolic functions. J. Agric. Sci., Mansoura Univ., 32(3):1029- 1037.
- Isaac, O. (1979). Pharmacological investigations with compounds of chamomile 1. On the pharmacology of (-)- $\alpha$ -bisabolol and bisabolol oxides (review). *Planta Med.*, 35:118-124.
- Jakobsen, P.E.; S.G. Kirsten and H. Niclsen (1960). Fredjelighed frogmed fierbrae. Digestibility trails with poultry "Bereting fra for sogslabortoriat, Kabenhaven, 56: 1-34.

- Khaki, A.; Fatemek, F.; Mohammad, N.; Amir, A.K.; Chelar, C.O.; Marefat, N. and Mohammad, H. (2009). The effects of ginger on spermatogenesis and sperm parameters. *Iranian J. Reproduc. Med.*, 7(1): 7-12.
- Khodary, R.M.; M.H.El-Ezzawy and I.R.Hamdy(1996). Effect of *Niegalia sativae* on egg production, hatchability percentage and some biochemical values in laying hens with reference to fertility in cockerles. *Proc. of 7<sup>th</sup> Sci. Vet. Med. Assuit Univ. Egypt.*, 17-19.
- Leung, A.Y.; and S. Foster, (1996). *Encyclopedia of Common Natural Ingredients used in Food, Drugs and Cosmetics*, 2nd ed. New York: John Wiley & Sons, Inc.
- Mann, C. and E.J.Staba,(1986). The chemistry, pharmacology and formulations of Chamomile . In, Graker L.E., Simon J. E., eds. *Herbs, spices and Medicinal plants: Recent Advances in Bolany, Horticulture and pharmacology* vol. 1. Arizona, Oryxprss, 235-280.
- McKay, D.L. and J.B .Blumberg (2009). A review of the bioactivity and potential health benefits of chamomile tea (*Matricaria recutita* L.). Human Nutrition Research Center on Aging at Tufts University, USA. 20(7):519- 530.
- Melrose, D. R. and Laing, J. A. (1970). The characteristics of normal semen. Chap. 4-*Fertility and infertility in the Domestic animals*. Ed. by J. A. Laing Bailliere Tindall and Gasssell, London.
- Morsy A.S, (2007). Enhancing rabbit performance reared in newly reclaimed areas by alleviating the environmental stress. Ph.D. Thesis, Fac. Agric., Cairo Univ., Egypt.
- Nadia, L.R. (2003). Effect of using some medicinal plants on performance and immunity of broiler chicks. Thesis, Fac. of Agric., Cairo Univ., Egypt.
- Namur, A. P.; J Morel and H. Bichek, (1988). Compound animal feed and feed additives. In Deboer, F., H. BICHEI, eds. *Livestock feed resources and feed evaluation in Europe*. Elsevier. Sci. publ., Amsterdam.
- N.R.C. (1994). National Research Council, Nutrient requirements of poultry 9<sup>th</sup> ed. National Academic of Science. Washington, Dc. USA.
- Osman, S.M.H. (1996). Responses of broiler chicks for some dietary supplementation M.Sc. Thesis Fac., of Agric. Zagazig Univ., Egypt.
- Riyad (1987). Effect of fenugreek and lupine seeds on the development of experimental diabetes in rats. *Planta Medica* 286-289.
- SAS (1996). *Statistical analysis system User's Guide*. SAS institute, Gary, North Carolina.
- Schelicher, A.; Fritz Z. and Kinal S. (1998). The use of some herbs in concentrates for broiler chickens. *Rocz. Nauk. Zoot*, 25 (3): 213-224.
- Sedki, A.A. (1999). Effect of different housing systems on rabbits performance. Ph.D. Thesis, Fac. of Agric., Cairo Univ., Egypt.
- Sharma, (1986). Effect of fenugreek seeds and leaves on blood glucose and serum insulin responses in human subjects. *Nutrition Research* 6: 1353-1364.
- Smith, J. T. and Mayer, D. T. (1955). Evaluation of sperm concentration by the haemocytometer method. *Fertil. Steril.*, 6: 271-275.

- Titus, H.W. and J.C. Fritz(1971). The scientific feeding of chickens. The interstate printers and Publishers, Inc., Danville, Illinois, U.S.A.
- White, I. G. (1976). Reproduction in the male. In: *Veterinary Physiology*. Phillis, J.G. Bristol: Wright-Scientechnica.
- Zeid, A. M. (1998). Effect of medicinal herbs in performance of goat. Ph. D. Faculty of Agric. Cairo University.

### تأثير مسحوق ازهار البابونج كأضافات غذائية فى علائق السمان الياباني البياض على الاداء الانتاجى والتناسلى

خالد عبد الجليل\*، هنده عبد الجليل محمود\*، أمال محمد حسن\*\* و على صابر مرسى\*\*

\* قسم تغذية الحيوان والدواجن - مركز بحوث الصحراء - المطرية- القاهرة.  
\*\* قسم فسيولوجيا الحيوان والدواجن - مركز بحوث الصحراء - المطرية- القاهرة.

استخدم في هذا البحث عدد ١٨٠ سمان ياباني ( ١٢٠ أنثى و ٦٠ ذكر) من عمر ٤ أسابيع وحتى عمر ٢٢ أسبوع. قسمت الإناث عشوائيا الى اربع مجموعات تجريبية متساوية (٣٠ أنثى / مجموعة) و قسمت الذكور ( ٦٠ ذكر) إلى أربع مجاميع تجريبية متساوية ( تم حفظ كل ذكر بطريقة فرديه ).

غذيت المجموعة الاولى على عليقة مقارنه بينما غذيت الثلاثه مجاميع الاخرى على عليقة المقارنه مضاف اليها نسب متدرجه من مسحوق ازهار البابونج ٠.٢٥ و ٠.٥٠ و ٠.٧٥ جم/كجم عليقه. العلائق المستخدمه متشابهه في نسبة البروتين الخام ٢٠% والطاقة الممتله ٢٩٠٠ كيلو كالورى /كيلوجرام.

الهدف من البحث دراسه تأثير اضافة مسحوق ازهار البابونج (كنبات طبي) إلى عليقه السمان الياباني البياض على الاداء الانتاجى والتناسلى.

#### ويمكن إيجاز أهم النتائج في النقاط التاليه:

- سجل وزن الجسم والتغير في وزن الجسم خلال الفترة التجريبية تحسن معنوي ( عند المستوى ٥%) بزيادة مستوى إضافة مسحوق أزهار البابونج، حيث سجلت المعامله المغذاه على ٠.٧٥ جم /كجم تليها المعامله المغذاه على ٠.٥٠ جم /كجم من إضافة مسحوق أزهار البابونج تحسنا في كل من وزن الجسم والتغير في وزن الجسم، بينما سجلت معاملة المقارنه اكثر القيم انخفاضاً.
- لوحظ زيادة معدل استهلاك الغذاء خلال فتره التجربه زيادة معنويه (عند مستوى ٥%) وذلك بإضافة مسحوق ازهار البابونج في العليقه، وقد سجلت المعامله المغذاه على ٠.٧٥ جم /كجم أعلى تلك القيم، بينما سجلت معاملة المقارنه اقل القيم خلال الفتره التجريبية.
- حققت المعامله التي غذيت على ٠.٥٠ جم /كجم من اضافة مسحوق ازهار البابونج افضل معدل تحويل غذائي خلال فترة التجربه مقارنه بباقي المعاملات.
- لم يسجل العمر عند النضج الجنسي ووزن البيضة الأولى فروق معنويه، بينما سجل وزن وعدد وكتلة البيض اختلاف معنويا (عند مستوى ٥%) بين المعاملات أثناء الفتره التجريبية.
- سجلت كلامن النسبه المئويه للفقس والخصوبه فروق معنويه ( عند مستوى ٥%) بين مجموعات التجريبية.
- سجلت النسبه المئويه للصفار وسمك القشره زياده غير معنويه وذلك بزيادة مستوى الإضافة لمسحوق أزهار البابونج في العليقه.
- سجلت النسبه المئويه للبياض و دليل الصفار انخفاض معنويا ( عند مستوى ٥%) بينما سجلت القشره وشكل البيضة زياده معنويه ( عند مستوى ٥%) وذلك بزيادة مستوى الإضافة لمسحوق أزهار البابونج في العليقه.

- أظهرت معاملات الهضم الظاهريه للماده العضويه والبروتين الخام والألياف الخام و مستخلص الأثير والمستخلص الخالي من النتروجين والقيم الغذائية اختلافا معنوياً ( عند مستوى ٥%) بإضافه مسحوق ازهار البابونج في العليقه.
  - تحقق أعلى عائد اقتصادي للمعامله المغذاه على ٠.٥٠ جم/كجم من مسحوق ازهار البابونج خلال فترة التجربه، مقارنة بباقي المعاملات بينما سجلت معاملة المقارنه اقل القيم.
  - لم يكن هناك اختلافات معنويه فى كل من حجم القذفه وعدد الحيوانات المنويه الكليه ونسبة الحيوانات المنويه المشوهه بين المجموعات المعامله.
  - حققت المعامله التي غذيت على ٠.٥٠ جم /كجم من اضافه مسحوق ازهار البابونج تحسنا معنوياً ( عند مستوى ٥%) فى تركيز الحيوانات المنويه وصفات حركة الحيوانات المنويه والحركة الكليه للحيوانات المنويه ونسبة الحيوانات المنويه الحيه ودليل جودة السائل المنوى.
  - اضافة مسحوق ازهار البابونج (٠.٥ جم/كجم) خفضت معنوياً ( عند مستوى ٥ %) رقم الأس الهيدروجيني بحوالى ٨.٥ % مقارنة بمجموعة المقارنه.
  - حققت المعامله التي غذيت على ٠.٥٠ جم /كجم من اضافة مسحوق ازهار البابونج زياده معنويه ( عند مستوى ٥%) فى الهيموجلوبين والبروتين الكلى والجلوبيولين،
  - انخفض معنوياً تركيز الكولستيرول فى المعامله التي غذيت على ٠.٥٠ و ٠.٧٥ جم /كجم.
  - لم يكن هناك اختلافات معنويه فى كل من تركيزات الجلوكوز والألبومين والكرياتينين وأنزيم ألانين ترانس ٨ أمينيز وأنزيم أسبارتات ترانس أمينيز.
- يمكن التوصيه من الوجهه الغذائيه والفسولوجيه والاقتصاديه إلى امكانيه استخدام مسحوق ازهار البابونج في العليقه بمستوى ٠.٥٠ جم/كجم في علائق السمان البياض لتحسين الاداء دون تأثير سلبي على معدلات الإنتاج والعائد الاقتصادي وبعض الوظائف الفسيولوجية.

#### قام بتحكيم البحث

كلية الزراعة – جامعة المنصورة  
مركز بحوث الصحراء

أ.د / أمينة عبد المطلب السروي  
أ.د / رافت السيد خضير